

Neuronal Excitability

Differential Impact of Inhibitory G-Protein Signaling Pathways in Ventral Tegmental Area Dopamine Neurons on Behavioral Sensitivity to Cocaine and Morphine

Margot C. DeBaker,¹  Ezequiel Marron Fernandez de Velasco,² Nora M. McCall,¹ Anna M. Lee,² and  Kevin Wickman²

<https://doi.org/10.1523/ENEURO.0081-21.2021>

¹Graduate Program in Neuroscience, University of Minnesota, Minneapolis, MN 55455 and ²Department of Pharmacology, University of Minnesota, Minneapolis, MN 55455

Abstract

Drugs of abuse engage overlapping but distinct molecular and cellular mechanisms to enhance dopamine (DA) signaling in the mesocorticolimbic circuitry. DA neurons of the ventral tegmental area (VTA) are key substrates of drugs of abuse and have been implicated in addiction-related behaviors. Enhanced VTA DA neurotransmission evoked by drugs of abuse can engage inhibitory G-protein-dependent feedback pathways, mediated by GABA_B receptors (GABA_BRs) and D₂ DA receptors (D₂Rs). Chemogenetic inhibition of VTA DA neurons potently suppressed baseline motor activity, as well as the motor-stimulatory effect of cocaine and morphine, confirming the critical influence of VTA DA neurons and inhibitory G-protein signaling in these neurons on this addiction-related behavior. To resolve the relative influence of GABA_BR-dependent and D₂R-dependent signaling pathways in VTA DA neurons on behavioral sensitivity to drugs of abuse, we developed a neuron-specific viral CRISPR/Cas9 approach to ablate D₂R and GABA_BR in VTA DA neurons. Ablation of GABA_BR or D₂R did not impact baseline physiological properties or excitability of VTA DA neurons, but it did preclude the direct somatodendritic inhibitory influence of GABA_BR or D₂R activation. D₂R ablation potentiated the motor-stimulatory effect of cocaine in male and female mice, whereas GABA_BR ablation selectively potentiated cocaine-induced activity in male subjects only. Neither D₂R nor GABA_BR ablation impacted morphine-induced motor activity. Collectively, our data show that cocaine and morphine differ in the extent to which they engage inhibitory G-protein-dependent feedback pathways in VTA DA neurons and highlight key sex differences that may impact susceptibility to various facets of addiction.

Key words: cocaine; CRISPR; D₂ dopamine receptor; GABA_B receptor; morphine; sex differences

Significance Statement

Although inhibitory G-protein-dependent signaling involving the GABA_B receptor (GABA_BR) and D₂ dopamine (DA) receptor (D₂R) in ventral tegmental area (VTA) DA neurons is thought to limit DA neurotransmission evoked by drugs of abuse, their relative impact on behavioral sensitivity to such drugs is unclear. Using a neuron-specific viral CRISPR/Cas9 approach, we show that loss of D₂R in VTA DA neurons enhances behavioral sensitivity to systemic administration of cocaine in male and female mice, whereas loss of GABA_BR enhances cocaine sensitivity only in males. Neither GABA_BR nor D₂R ablation impacted behavioral sensitivity to morphine. Thus, differential engagement of inhibitory feedback pathways in VTA DA neurons likely contributes to drug-specific neurophysiological and behavioral effects and may underlie sex differences associated with some facets of addiction.

Received March 2, 2021; accepted March 3, 2021; First published March 10, 2021.

The authors declare no competing financial interests.

Author contributions: K.W., M.C.D., E.M.F.d.V., N.M.M., and A.M.L. designed research; M.C.D. and N.M.M. performed research; E.M.F.d.V. contributed unpublished reagents/analytic tools; M.C.D. and N.M.M. analyzed data; K.W., M.C.D., and A.M.L. wrote the paper.

Introduction

Dopamine (DA) neurons of the ventral tegmental area (VTA) are an integral part of the mesocorticolimbic system, a network of brain regions that mediates responses to natural rewards and drugs of abuse (Lammel et al., 2014; Volkow and Morales, 2015; Juarez and Han, 2016; Lüscher, 2016; Schultz, 2016). Drugs of abuse enhance DA neurotransmission in the mesocorticolimbic system via actions on distinct molecular targets (Di Chiara and Imperato, 1988; Lüscher and Ungless, 2006). Cocaine, for example, inhibits monoamine transporters, including the DA transporter (DAT), which recycles DA from the extracellular space, allowing DA levels to rise and persist in the VTA and its projection targets (Groves et al., 1975; Beart et al., 1979; Di Chiara and Imperato, 1988; Chen et al., 1996; Iravani et al., 1996; Adell and Artigas, 2004; Kita et al., 2009; Adrover et al., 2014). In contrast, morphine suppresses inhibitory GABAergic input, enhancing VTA DA neuron activity via disinhibition (Johnson and North, 1992; Zhou et al., 2009; Jalabert et al., 2011).

While enhanced DA neurotransmission underlies addiction-related behaviors (Hyman et al., 2006), it can also trigger negative feedback mediated by inhibitory G-protein signaling pathways (Steketee and Kalivas, 1990, 1991; Narayanan et al., 1996; Rahman and McBride, 2000). For example, cocaine elevates DA levels in the VTA and terminal regions, provoking feedback inhibition of VTA DA neurons via activation of D₂ DA receptors (D₂Rs) in the somatodendritic compartment and axon terminals (Einhorn et al., 1988; Bradberry and Roth, 1989; Brodie and Dunwiddie, 1990; Beckstead et al., 2004; Ford, 2014). Cocaine also provokes GABAergic feedback to VTA DA neurons indirectly, by stimulating D₁R-expressing GABAergic medium spiny neurons in the nucleus accumbens (NAc) that project to the VTA (Rahman and McBride, 2001; Watabe-Uchida et al., 2012; Menegas et al., 2015; Edwards et al., 2017; Pignatelli and Bonci, 2018). This “long-loop” GABAergic feedback to VTA DA neurons is mediated primarily by activation of somatodendritic GABA_B receptors (GABA_BRs) on VTA DA neurons (Edwards et al., 2017).

Pharmacological and genetic approaches have implicated GABA_BR and D₂R-dependent signaling in the VTA in drug-induced behaviors. For example, pharmacological activation of D₂R in the VTA suppressed cocaine-induced activity (Koulchitsky et al., 2016) and cocaine-reinstated

drug seeking behavior (Xue et al., 2011), while pharmacological inhibition of D₂R in the VTA increased psychostimulant-induced locomotor activity (Chen and Reith, 1994; Tanabe et al., 2004). Similarly, intra-VTA infusion of the GABA_BR agonist baclofen blocked the locomotor-stimulatory effect of psychostimulants and opioids (Kalivas et al., 1990; Steketee and Kalivas, 1991; Chen and Reith, 1994; Leite-Morris et al., 2002, 2004) and attenuated self-administration of cocaine, opioids, and other drugs of abuse (Xi and Stein, 1999; Brebner et al., 2000; Leite-Morris et al., 2004; Backes and Hemby, 2008). RNAi-mediated suppression of D₂R in the rat VTA (cell non-selective) enhanced cocaine-related behavior (de Jong et al., 2015; Chen et al., 2018), enhanced choice impulsivity (Bernosky-Smith et al., 2018), and increased functional brain activity (Martin et al., 2020). Ablation of D₂R in DA neurons throughout the mouse brain correlated with enhanced cocaine-induced activity (Bello et al., 2011), as well as acquisition of cocaine self-administration and reactivity to drug-paired cues (Holroyd et al., 2015). Interestingly, partial suppression of GABA_BR in VTA DA neurons unmasked cocaine-induced activity normally absent in BALB/c mice, but did not impact morphine-induced activity (Edwards et al., 2017).

Collectively, available data suggests that D₂R and GABA_BR-dependent signaling pathways in VTA DA neurons may exert a differential influence on behavioral sensitivity to drugs of abuse. Published studies investigating D₂R and GABA_BR signaling in VTA DA neurons, however, have used approaches that provide either anatomic or cellular specificity, or do not completely suppress inhibitory signaling, or focus only on one signaling pathway or drug of abuse. Accordingly, the goal of this study was to compare the impact of D₂R or GABA_BR ablation in VTA DA neurons on behavioral sensitivity to cocaine and morphine. To this end, we developed a neuron-specific viral CRISPR/Cas9 approach to ablate D₂R and GABA_BR selectively in VTA DA neurons of adult mice. Our findings show that D₂R and GABA_BR-dependent signaling exert drug-specific and sex-specific influences on motor activity in mice.

Materials and Methods

Animals

All studies were approved by the Institutional Animal Care and Use Committee at the University of Minnesota. The B6.SJL-*Slc6a3*^{tm1.1(cre)Bkmn}/J (stock #006660, The Jackson Laboratory) knock-in line was used in this study; heterozygous subjects, referred to throughout as DATCre (+) mice, were generated by crossing with C57BL/6J subjects. DATCre(+) mice were also crossed in multiple rounds with a Cre-dependent Cas9GFP knock-in line (B6;129-Gt(ROSA)26Sor^{tm1(CAG-cas9x,-EGFP)Fez}/J, The Jackson Laboratory, stock #026179), to generate DATCre(+) subjects homozygous for the Cas9GFP(+) mutation; these mice are referred to throughout as DATCre(+);Cas9GFP(+) mice. All mice used in experiments were bred in-house. Mice were group housed, maintained on a 14/10 h light/dark cycle and were provided *ad libitum* access to food and water.

This work was supported by National Institutes of Health Grants DA034696 and AA027544 (to K.W.), DA007234 (to M.C.D.), DA041767 (to N.M.M.), AA026598 (to A.M.L.), and the University of Minnesota Viral Vector and Cloning Core (DA048742).

Acknowledgements: We thank Hannah Oberle, Mehrsa Zahiremani, and Courtney Wright for exceptional care of the mouse colony and the Genome Engineering and iPSC Center of Washington University for designing and validating gRNA sequences used in this study.

Correspondence should be addressed to Kevin Wickman at wickm002@umn.edu.

<https://doi.org/10.1523/ENEURO.0081-21.2021>

Copyright © 2021 DeBaker et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license, which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Reagents

Baclofen, quinpirole, and sulpiride were purchased from Sigma, and CGP54626 and clozapine-N-oxide (CNO) were purchased from Tocris. Cocaine and morphine were obtained through Boynton Health Pharmacy at the University of Minnesota. All adeno-associated viruses (AAVs) were packaged in AAV8 serotype by the University of Minnesota Viral Vector and Cloning Core (Minneapolis, MN) following standard packaging procedures (Chen et al., 2019), titers were between 0.2 and 4×10^{14} genocopies/ml. The packaging plasmids (pRC8 and pHelper) were obtained from the University of Pennsylvania Vector Core. The plasmids pAAV-hSyn-hM4Di(mCherry) and pAAV-hSyn-DIO-mCherry (Addgene plasmids #50475 and #50459, respectively) were gifts from Bryan Roth. To obtain the pAAV-hSyn-DIO-hM4Di(mCherry) plasmid, the mCherry from the pAAV-hSyn-DIO-mCherry was replaced by an hM4Di(mCherry) cassette via subcloning. For the CRISPR/Cas9 experiments the pAAV-U6-gRNA-hSyn-NLSmCherry was generated using the backbone of the plasmid pAAV-U6-gRNA-hSyn-Cre-2A-EGFP-KASH (Platt et al., 2014; Addgene plasmid #60231) that was a gift from Feng Zhang. The Genome Engineering and iPSC Center of Washington University designed and tested guide RNA (gRNA) sequences targeting the *Drd2* (*D₂R*) and *Gabbr1* (*GABA_BR1*) genes. Sequences used were as follows: *D₂R*, CATGACAGTAACTCGGCGCT; *GABA_BR1*, ACGGCGTGAGTATACATCG; *LacZ*, TGCGAATACGCC CACGCGAT.

Intracranial manipulations

Mice (>45 d) were placed in a stereotaxic frame (David Kopf Instruments) under isoflurane anesthesia. Microinjectors, made by affixing a 33-gauge stainless steel hypodermic tube within a shorter 26-gauge stainless steel hypodermic tube, were attached to polyethylene-20 tubing affixed to 10- μ l Hamilton syringes, and were lowered through burr holes in the skull to the VTA (from bregma: -2.75 mm A/P, ± 0.55 – 0.7 mm M/L, -5 mm D/V); 300–500 nl of virus was injected per side at a rate of 100 nl/min. The optimized coordinates and viral load ensured full coverage of the VTA along anterior/posterior and rostral-caudal axes, with minimal spread into the adjacent substantia nigra pars compacta. Microinjectors were left in place for 10 min following infusion to reduce solution backflow along the infusion track. Slice electrophysiology and behavioral experiments were performed three to four and five to six weeks after viral infusion for chemogenetic and CRISPR experiments, respectively.

Slice electrophysiology

Electrophysiological properties of VTA DA neurons were evaluated in behaviorally naive adult mice (66–93 d). Horizontal slices (225 μ m) containing the VTA were prepared in ice-cold sucrose substituted ACSF, and allowed to recover at room temperature in ACSF containing the following: 125 mM NaCl, 2.5 mM KCl, 1.25 mM NaH_2PO_4 , 25 mM NaHCO_3 , 11 mM glucose, 1 mM MgCl_2 , and 2 mM CaCl_2 , pH 7.4, for at least 1 h, as described (McCall et al.,

2017). Neurons in the lateral VTA exhibiting appropriate fluorescence were targeted for analysis as this sub-region of the VTA receives prominent input from the NAc that mediates *GABA_BR*-dependent feedback (Edwards et al., 2017). Whole-cell data were acquired using a Multiclamp 700A amplifier and pCLAMPv.9.2 software (Molecular Devices, LLC), using recording conditions described in previous publications (McCall et al., 2017). Input/membrane resistance (R_M) and apparent capacitance (C_M) were determined using a 5 mV/10 ms voltage step, with current responses filtered at 5 Hz. Immediately after establishing whole-cell access, I_h conductance was measured using a 200-ms voltage step to -120 mV; the difference in current from beginning to end of the -120 -mV step was taken as I_h amplitude. Subsequently, spontaneous activity was measured in current-clamp mode ($I = 0$) for 1 min. Neurons exhibiting no spontaneous activity were not evaluated. Action potential half-width (AP_{50}) was determined by averaging five AP_{50} values. For rheobase assessments, cells were held in current-clamp mode at -80 pA to prevent spontaneous activity, and then given 1-s current pulses, beginning at -100 pA and progressing in 20-pA increments. Rheobase was defined as the minimum current step that evoked one or more action potentials. In chemogenetic experiments, the change in rheobase measured before and after bath application of CNO application was determined. Somatodendritic holding currents were measured in voltage-clamp mode ($V_{\text{hold}} = -60$ mV) following bath application of CNO (10 μ M), baclofen (200 μ M), or quinpirole (10 μ M). All command potentials factored in a junction potential of -15 mV predicted using JPCalc software (Molecular Devices, LLC). Series and input resistances were tracked throughout the experiment. If series resistance was unstable or exceeded 20 M Ω , the experiment was excluded from analysis.

Locomotor activity

Locomotor activity was assessed in open-field activity chambers housed in sound-attenuating cubicles (Med-Associates). Each cubicle was equipped with three 16-beam infrared arrays permitting automated measurements of distance traveled (Activity Monitor 5; Med-Associates). Animals were habituated to the testing room for at least 30 min before testing. Subjects were acclimated over 3 d; on day 1, animals were handled and placed in the open field for 60 min. On days 2 and 3, animals were given an intraperitoneal injection of saline and placed in the open field for 60 min. For DREADD experiments, mice were injected with CNO (2 mg/kg, i.p.) 30 min before either saline, cocaine (15 mg/kg, i.p.), or morphine (10 mg/kg, i.p.) injection on day 4. Distance traveled during the 60-min period following saline or drug injection was measured; separate cohorts of mice received saline, cocaine and morphine injections. For CRISPR/Cas9 experiments, mice were placed in the open field for 30 min before injection each day to acclimate to the chamber. Activity was measured on day 3 following saline injection, and again on day 4 following injection of cocaine (15 mg/kg, i.p.) or morphine (10 mg/kg, i.p.), separate cohorts of mice were used for cocaine and morphine studies.

Thigmotaxis was quantified by dividing the distance traveled in the periphery by the total distance traveled, as described (Pravetoni and Wickman, 2008). After behavioral testing, the scope and accuracy of viral targeting was assessed by fluorescence microscopy; 225- μm horizontal slices of the midbrain were obtained using a vibratome and images were acquired on an Olympus IX-80 microscope using MetaMorph Advanced Acquisition v.7.7.7.0 software (Molecular Devices, LLC). Only data from animals with bilateral viral-driven fluorescence, where the majority of fluorescence was confined to VTA (with minimal spread to the adjacent substantia nigra), were included in the final analysis.

Statistical analysis

Data are presented throughout as the mean \pm SEM. Statistical analyses were performed using Prism v.9 software (GraphPad Software). All studies included balanced numbers of male and female mice, and data were analyzed first for sex effects. If no sex differences were observed, data from male and female subjects were pooled. If any data point fell outside the range of 2 SDs from the mean, it was excluded as an outlier. Across the entire study, this outlier detection approach led to the removal of one point from the hM4Di/morphine activity dataset and 1 point from the hM4Di control/morphine activity dataset. Differences were considered significant if $p < 0.05$.

Results

G-protein-dependent inhibition of VTA DA neurons suppresses motor activity

Motor activation is an unconditioned DA-dependent response in mice to systemic administration of cocaine and morphine (van Rossum et al., 1962; Delfs et al., 1990; Kalivas and Duffy, 1993) that can be recapitulated by direct chemogenetic or optogenetic stimulation of VTA DA neurons (Kim et al., 2012; Tye et al., 2013; Boender et al., 2014; Guo et al., 2014). To test whether inhibitory G-protein-dependent signaling in VTA DA neurons can suppress the locomotor stimulatory effect of systemic cocaine and morphine, we used DATCre(+) mice and a Cre-dependent viral inhibitory chemogenetic approach. Cre-dependent AAV vectors harboring either hM4Di (mCherry) or mCherry control were infused into the VTA of adult male and female DATCre(+) mice to permit selective chemogenetic inhibition of VTA DA neurons (Fig. 1A).

In acutely isolated midbrain slices from virally treated DATCre(+) mice, bath application of CNO decreased the excitability (increased the rheobase) of hM4Di-expressing, but not mCherry control, DA neurons (Fig. 1B,C). CNO also evoked a somatodendritic inhibitory current ($V_{\text{hold}} = -60\text{ mV}$) reversed by 0.3 mM Ba^{2+} in hM4Di (mCherry)-expressing but not control VTA DA neurons (Fig. 1D,E). Thus, chemogenetic inhibition of VTA DA neurons, like the direct somatodendritic inhibition evoked by D_2R and $\text{GABA}_{\text{B}}\text{R}$ activation, is likely mediated by activation of G-protein-gated inwardly rectifying K^+ (GIRK/Kir3) channels (Beckstead et al., 2004; Cruz et al., 2004; Labouèbe et al., 2007; Arora et al., 2010).

Consistent with a previous report (Runegaard et al., 2018), chemogenetic inhibition of VTA DA neurons suppressed motor activity evoked by injection of saline, as well as cocaine and morphine (Fig. 1F). These results confirm that VTA DA neurons are key regulators of motor activity in mice and show that activation of inhibitory G-protein signaling in these neurons can potently suppress baseline activity and the motor-stimulatory effects of cocaine and morphine.

CRISPR/Cas9 ablation of D_2R and $\text{GABA}_{\text{B}}\text{R}$ in VTA DA neurons

To assess the impact of D_2R and $\text{GABA}_{\text{B}}\text{R}$ -dependent signaling pathways in VTA DA neurons on behavioral sensitivity to cocaine and morphine, we developed a DA neuron-specific, viral CRISPR/Cas9 ablation approach. DATCre(+) mice were crossed with a Cre-dependent Cas9GFP line to generate DATCre(+):Cas9GFP(+) mice. AAV vectors harboring gRNAs targeting LacZ (control), D_2R , or $\text{GABA}_{\text{B}}\text{R1}$ were generated and infused into the VTA of male and female DATCre(+):Cas9GFP(+) mice (Fig. 2A).

To assess the efficacy and selectivity of the viral vectors, somatodendritic currents evoked by bath application of the $\text{D}_{2/3}\text{R}$ agonist quinpirole and the $\text{GABA}_{\text{B}}\text{R}$ agonist baclofen were measured in VTA DA neurons five to six weeks after viral infusion. In slices from LacZ gRNA-treated subjects, somatodendritic outward inhibitory currents were reliably evoked by quinpirole and baclofen (Fig. 2B), consistent with previous studies (Beckstead et al., 2004; Cruz et al., 2004; Labouèbe et al., 2007; Arora et al., 2010; McCall et al., 2017). In VTA DA neurons from mice treated with D_2R gRNA, quinpirole-induced currents were completely absent (Fig. 2B,C), while baclofen (applied after quinpirole) evoked normal responses (Fig. 2B,D). Interestingly, a sex difference in the amplitude of quinpirole-induced responses was observed, with currents in VTA DA neurons from female subjects larger than those in males (Fig. 2C). No sex differences were observed in the amplitude of baclofen-induced currents.

In VTA DA neurons from mice treated with $\text{GABA}_{\text{B}}\text{R1}$ gRNA, somatodendritic responses to baclofen were completely absent (Fig. 2E,F). Responses to quinpirole (applied after baclofen application) were larger than those seen in VTA DA neurons from LacZ-treated controls (Fig. 2G). As this difference may reflect the impact of prior activation by $\text{GABA}_{\text{B}}\text{R}$ of a shared effector in control cells, we conducted a separate study where only quinpirole was applied to the slice. In these experiments, quinpirole-induced somatodendritic currents were not significantly different between $\text{GABA}_{\text{B}}\text{R}$ -lacking and control VTA DA neurons (Fig. 2H).

We also examined the impact of D_2R and $\text{GABA}_{\text{B}}\text{R}$ ablation on VTA DA neuron excitability, as measured by spontaneous activity and rheobase. No sex differences were observed with respect to either excitability measure and, as such, data from male and female subjects were pooled to increase the power of the study. Ablation of D_2R or $\text{GABA}_{\text{B}}\text{R}$ did not impact spontaneous activity of VTA DA neurons (Fig. 2I) or rheobase (Fig. 2J). Similarly,




Figure 1. Inhibitory G-protein signaling in VTA DA neurons suppresses baseline and drug-induced motor activity. **A**, Viral targeting in a DATCre(+) mouse treated intra-VTA AAV8-hSyn-DIO-mCherry, with rostral-caudal tiling of panels highlighting viral spread (mCherry fluorescence). ml, medial lemniscus; MT, medial terminal nucleus of the accessory optic tract; PBP, parabrachial pigmented nucleus of the VTA; SNc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata; VTAR, rostral part of the VTA. Scale bar: 500 μ m. **B**, Example of rheobase measurement in a VTA DA neuron expressing hM4Di, before and after bath perfusion of CNO (10 μ M). The traces shown were the first to display spiking and were recorded following injection of the current indicated in red below. **C**, Change in rheobase induced by CNO (10 μ M) in VTA DA neurons from DATCre(+) mice treated with AAV8-hSyn-DIO-hM4Di(mCherry) or AAV8-hSyn-DIO-mCherry control ($t_{(18)} = 4.555$; $***p = 0.0002$; unpaired Student's t test; $n = 9$ –11/group). **D**, Somatodendritic inhibitory currents ($V_{\text{hold}} = -60$ mV) evoked by CNO (10 μ M) in VTA DA neurons from DATCre(+) mice treated with AAV8-hSyn-DIO-hM4Di(mCherry) or AAV8-hSyn-DIO-mCherry control. **E**, Summary of CNO-induced somatodendritic currents in VTA DA neurons from DATCre(+) treated with AAV8-hSyn-DIO-hM4Di(mCherry) or AAV8-hSyn-DIO-mCherry control ($t_{(16)} = 3.732$; $**p = 0.0018$; unpaired Student's t test; $n = 9$ /group). **F**, Total distance traveled in an open field during the 60-min interval following injection of saline ($t_{(16)} = 4.385$, $***p = 0.0005$), 15 mg/kg cocaine ($t_{(14)} = 11.20$, $***p < 0.0001$), and 10 mg/kg morphine ($t_{(15)} = 9.401$, $***p < 0.0001$) in DATCre(+) mice treated with intra-VTA AAV8-hSyn-DIO-hM4Di(mCherry) or AAV8-hSyn-DIO-mCherry control ($n = 8$ –9 mice/group). CNO (2 mg/kg, i.p.) was administered to all subjects 30 min before saline or drug challenge.

no impact of D₂R or GABA_BR ablation was detected on I_h current amplitude, cell capacitance, action potential half-width, or input resistance (Table 1). Collectively, these data show that we can selectively ablate GABA_BR-dependent or D₂R-dependent signaling in VTA DA neurons, and that loss of these signaling pathways does not impact baseline electrophysiological characteristics of VTA DA neurons

Impact of D₂R and GABA_BR in VTA DA neurons on baseline and cocaine-induced activity

We next examined the impact of D₂R or GABA_BR ablation on open-field motor activity measured after injection of saline or cocaine. Given the observed sex difference in the strength of D₂R-dependent signaling in VTA DA neurons, and our prior report of a sex difference in cocaine-induced motor activity in mice (McCall et al., 2017), we analyzed data from male and female subjects separately.

In support of this approach, we found that while male and female LacZ control mice showed no difference in total distance traveled after cocaine injection, the temporal profile of cocaine-induced activity was notably different for male and female subjects (Extended Data Fig. 3-1A), female subjects showed sharper time-to-peak and decay phases relative to male subjects.

In females, we observed main effects of drug and viral treatment, as well as a significant interaction between drug and viral treatment, on total distance traveled in the postinjection interval. D₂R or GABA_BR ablation did not impact saline-induced activity in females (Fig. 3A, left). Loss of D₂R, but not GABA_BR, yielded enhanced cocaine-induced activity (Fig. 3A, right). Although the temporal profile of motor activity during the postinjection interval was qualitatively similar across the groups, loss of D₂R and GABA_BR in VTA DA neurons correlated with elevated activity levels at all time points following injection (Fig. 3B). Cocaine also enhanced thigmotaxis, as assessed by




Figure 2. Viral CRISPR/Cas9 ablation of GABA_B and D₂R in VTA DA neurons. **A**, Viral targeting in a DATCre(+):Cas9GFP(+) mouse treated with intra-VTA AAV8-U6-gRNA-hSyn-NLSmCherry; nucleus-localized mCherry fluorescence highlights the anatomic scope of viral targeting, and GFP fluorescence denotes the Cre-dependent expression of Cas9GFP in midbrain DA neurons of the VTA and substantia nigra. MT, medial terminal nucleus of the accessory optic tract; PBP, parabrachial pigmented nucleus of the VTA; SNc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata; VTAR, rostral part of the VTA. Scale bar: 500 μm. **B**, Somatodendritic inhibitory currents ($V_{\text{hold}} = -60$ mV) evoked by sequential bath application of quinpirole (quin; 20 μM) and then baclofen (bac; 200 μM) in VTA DA neurons from DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gD2R-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. Currents were reversed by the D_{2/3}R antagonist sulpiride (sulp; 5 μM) and GABA_BR antagonist CGP54626 (CGP; 2 μM). **C**, Summary of currents evoked by quinpirole (applied first) in VTA DA neurons from DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gD2R-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. Main effects of sex ($F_{(1,27)} = 6.391, p = 0.0176$) and viral treatment ($F_{(1,27)} = 104.6, p < 0.0001$) were detected, along with an interaction between sex and viral treatment ($F_{(1,27)} = 5.590, p = 0.0255$); # $p < 0.05$; *** $p < 0.001$ versus LacZ (within sex). **D**, Summary of currents evoked by baclofen, measured after quinpirole/sulpiride application (**C**), in VTA DA neurons from DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gD2R-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. There was no main effect of sex ($F_{(1,27)} = 1.741, p = 0.1981$) or viral treatment ($F_{(1,27)} = 1.183, p = 0.2864$), nor was there an interaction between sex and viral treatment ($F_{(1,27)} = 0.0047, p = 0.9461$). **E**, Somatodendritic inhibitory currents ($V_{\text{hold}} = -60$ mV) evoked by sequential bath application of baclofen (bac; 200 μM) and then quinpirole (quin; 20 μM) in VTA DA neurons from DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. Currents were reversed by CGP54626 (CGP; 2 μM) and sulpiride (sulp; 5 μM). **F**, Summary of currents evoked by baclofen (applied first) in VTA DA neurons from DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. A main effect of viral treatment was observed ($F_{(1,27)} = 120.9, p < 0.0001$), but there was no main effect of sex ($F_{(1,27)} = 0.1910, p = 0.6655$) or interaction between sex and viral treatment ($F_{(1,27)} = 0.2216, p = 0.6416$); +++ $p < 0.001$ (main effect of viral treatment). **G**, Summary of currents evoked by quinpirole, measured after baclofen/CGP54626 application (**F**) in VTA DA neurons from DATCre(+):Cas9GFP(+) mice treated

continued

with AAV8-U6-gGBR1-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. A main effect of viral treatment was observed ($F_{(1,24)} = 8.182, p = 0.0086$), but there was no main effect of sex ($F_{(1,24)} = 3.429, p = 0.0764$) or interaction between sex and viral treatment ($F_{(1,24)} = 1.970, p = 0.1733$); $++p < 0.001$ (main effect of viral treatment). **H**, Summary of currents evoked by quinpirrole in VTA DA neurons from a separate cohort of DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry or AAV8-U6-gLacZ-hSyn-NLSmCherry control. A main effect of sex was observed ($F_{(1,26)} = 13.41, p = 0.0011$), but there was no main effect of viral treatment ($F_{(1,26)} = 1.766, p = 0.1954$) or interaction between sex and viral treatment ($F_{(1,26)} = 0.0129, p = 0.9105$); $++p < 0.01$ (main effect of sex). **I**, Impact of D₂R or GABA_BR ablation on spontaneous activity in VTA DA neurons. No main effect of sex ($F_{(1,56)} = 0.7079, p = 0.4037$) or viral treatment ($F_{(2,56)} = 0.6571, p = 0.5223$) was detected, nor was there an interaction between sex and viral treatment ($F_{(2,56)} = 1.276, p = 0.2870$). **J**, Impact of D₂R or GABA_BR ablation on rheobase in VTA DA neurons. No main effect of sex ($F_{(1,56)} = 0.8080, p = 0.3725$) or viral treatment ($F_{(2,56)} = 1.495, p = 0.2332$) was detected, nor was there an interaction between sex and viral treatment ($F_{(2,56)} = 0.1573, p = 0.8548$).

calculating the ratio of distance traveled in the field periphery to total distance traveled (Fig. 3C). There was, however, no impact of D₂R or GABA_BR ablation in VTA DA neurons on thigmotaxis index (Fig. 3C).

In males as in females, we observed main effects of drug and viral treatment, and a significant interaction between drug and viral treatment on total distance traveled. Loss of D₂R or GABA_BR did not impact saline-induced activity (Fig. 3D, left). In contrast to our observations in females, however, ablation of D₂R or GABA_BR in males yielded comparably enhanced cocaine-induced activity over the 60-min postinjection interval (Fig. 3D, right). Interestingly, loss of D₂R in males correlated with higher levels of activity seen shortly after cocaine injection (Fig. 3E), yielding a temporal profile that was qualitatively similar to that observed in females (Extended Data Fig. 3-1B). As was the case with female subjects, thigmotaxis was significantly impacted by drug but not viral treatment (Fig. 3C). Thus, D₂R-dependent signaling in VTA DA neurons tempers behavioral sensitivity to cocaine in male and female mice, whereas GABA_BR-dependent signaling exerts an influence on behavioral sensitivity to cocaine in male mice only.

Impact of D₂R and GABA_BR in VTA DA neurons on baseline and morphine-induced activity

As we previously reported that the loss of GIRK channel activity in DA neurons in mice correlated with enhance motor stimulation in response to systemic administration of 10 mg/kg, morphine (Kotecki et al., 2015), we next compared the relative contribution of D₂R and GABA_BR to

this morphine-induced behavior. In female subjects, we observed a main effect of drug on motor activity, but no main effect of viral treatment or interaction between drug and viral treatment. Notably, no significant impact of D₂R or GABA_BR ablation on morphine-induced activity was observed in females (Fig. 4A, right). Although D₂R ablation correlated with elevated activity levels at all time points following morphine injection, activity levels were not significantly different from controls (Fig. 4B). Consistent with prior reports (Mickley et al., 1990), we observed a marked increase in thigmotaxis in all subjects after morphine injection, but there was no difference between viral treatment groups (Fig. 4C).

In males, we also observed a main effect of drug, but no main effect of viral treatment or interaction between drug and viral treatment. As was the case in females, loss of either D₂R or GABA_BR in males had no significant impact on morphine-induced activity during the 60-min interval (Fig. 4D, right) or on the temporal activity profile following morphine injection (Fig. 4E). We also observed a similar increase in thigmotaxis after injection of morphine as seen in females, with no difference between viral treatment groups (Fig. 4F). Thus, neither D₂R nor GABA_BR ablation in VTA DA neurons exerts a significant impact on morphine-induced motor activity in mice.

Discussion

Here, we used a neuron-specific viral CRISPR/Cas9 approach to compare the impact of D₂R or GABA_BR ablation in VTA DA neurons on baseline activity and behavioral sensitivity to cocaine and morphine. Loss of GABA_BR or

Table 1: Electrophysiological properties of VTA DA neurons

Sex	gRNA	N/n	C _M (pF)	R _M (MΩ)	I _n (pA)	AP ₅₀ (ms)
Male	LacZ	8/15	60 ± 2	200 ± 10	313 ± 49	0.83 ± 0.04
	GABA _B R1	3/8	56 ± 3	194 ± 11	240 ± 66	0.92 ± 0.07
	D ₂ R	3/8	62 ± 4	207 ± 34	246 ± 45	0.92 ± 0.04
			$F_{(2,28)} = 0.8299$ $p = 0.4465$	$F_{(2,28)} = 0.1024$ $p = 0.9030$	$F_{(2,28)} = 0.6179$ $p = 0.5463$	$F_{(2,28)} = 1.154$ $p = 0.3301$
Female	LacZ	9/16	62 ± 2	186 ± 11	274 ± 39	0.85 ± 0.04
	GABA _B R1	3/7	60 ± 4	203 ± 21	225 ± 70	0.89 ± 0.06
	D ₂ R	4/8	62 ± 3	174 ± 22	338 ± 71	0.99 ± 0.05
			$F_{(2,28)} = 0.2026$ $p = 0.8178$	$F_{(2,28)} = 0.6060$ $p = 0.5525$	$F_{(2,28)} = 0.7869$ $p = 0.4651$	$F_{(2,28)} = 2.431$ $p = 0.1063$

Data extracted from whole-cell recordings of VTA DA neurons from male and female DATCre(+):Cas9GFP(+) mice treated with intra-VTA control (LacZ), D₂R, or GABA_BR1 gRNA vectors. N/n, number of mice and individual experiments; C_M, apparent capacitance; R_M, input/membrane resistance; I_n, hyperpolarization-activated current; AP₅₀, action-potential half-width.




Figure 3. Impact of GABA_BR and D₂R ablation on the motor-stimulatory effect of cocaine. **A**, Total distance traveled during the 60-min period after injection of saline (left) or cocaine (15 mg/kg, i.p., right) in female DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry ($n = 9$), AAV8-U6-gD2R-hSyn-NLSmCherry ($n = 9$), or AAV8-U6-gLacZ-hSyn-NLSmCherry control ($n = 8$). Two-way repeated measures ANOVA revealed main effects of drug ($F_{(1,23)} = 328.2, p < 0.0001$) and viral treatment ($F_{(2,23)} = 3.778, p = 0.0381$), and an interaction between drug and viral treatment ($F_{(2,23)} = 4.600, p = 0.0209$); *** $p < 0.001$; ns, not significant ($p = 0.1209$). **B**, Distance traveled for female subjects before and after cocaine injection (denoted by arrow), evaluated in 5-min bins. Two-way repeated measures ANOVA revealed a main effect of bin ($F_{(3,055,70,26)} = 82.89, p < 0.0001$; Geisser–Greenhouse correction for sphericity) and viral treatment ($F_{(2,23)} = 4.613, p = 0.0207$), but there was no interaction between bin and viral treatment ($F_{(24,276)} = 0.9807, p = 0.4919$); + $p < 0.05$ (main effect of viral treatment). **C**, Thigmotaxis index for female subjects following saline or cocaine injection. Two-way repeated measures ANOVA revealed a main effect of drug ($F_{(1,23)} = 16.40, p = 0.0005$) but not viral treatment ($F_{(2,23)} = 0.07351, p = 0.9293$), and there was no interaction between drug and viral treatment ($F_{(2,23)} = 0.03955, p = 0.9613$); ++ $p < 0.001$ (main effect of drug). **D**, Total distance traveled during the 60-min period after injection of saline (left) or cocaine (15 mg/kg, i.p., right) in male DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry ($n = 8$), AAV8-U6-gD2R-hSyn-NLSmCherry ($n = 8$), or AAV8-U6-gLacZ-hSyn-NLSmCherry control ($n = 9$). Two-way repeated measures ANOVA revealed main effects of drug ($F_{(1,22)} = 91.27, p = 0.0157$) and viral treatment ($F_{(2,22)} = 4.173, p = 0.0291$), and an interaction between drug and viral treatment ($F_{(2,22)} = 5.050, p = 0.0157$); ** $p < 0.01$ and *** $p < 0.001$. **E**, Distance traveled for male subjects before and after cocaine injection (denoted by arrow), evaluated in 5-min bins. Two-way repeated measures ANOVA revealed a main effect of bin ($F_{(3,131,68,88)} = 21.10, p < 0.0001$) and viral treatment ($F_{(2,22)} = 4.747, p = 0.0193$), and an interaction between bin and viral treatment ($F_{(24,264)} = 4.743, p < 0.0001$); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (gD2R vs LacZ); # $p < 0.05$ and ## $p < 0.01$ (gGBR1 vs LacZ). **F**, Thigmotaxis index for male subjects following saline or cocaine injection. Two-way repeated measures ANOVA revealed a main effect of drug ($F_{(1,22)} = 21.75, p = 0.0001$), but no main effect of viral treatment ($F_{(2,22)} = 1.534, p = 0.2380$) or interaction between drug and viral treatment ($F_{(2,22)} = 0.9989, p = 0.3844$); ++ $p < 0.001$ (main effect of drug).

D₂R in VTA DA neurons had no significant impact on baseline activity, paralleling the lack of effect of these manipulations on excitability and other electrophysiological properties of VTA DA neurons. These behavioral findings are consistent with studies involving genetic suppression or ablation of D₂R or GABA_BR in the rodent VTA (de Jong et al., 2015; Edwards et al., 2017). Thus, D₂R-dependent and GABA_BR-dependent signaling pathways in VTA DA neurons exert minimal influence on baseline DA neuron excitability. Notably, mice lacking D₂R in DA neurons displayed hyperlocomotion (Bello et al., 2011), indicating that D₂R-dependent signaling in DA neuron populations outside of the VTA may regulate baseline motor activity.

D₂R ablation in VTA DA neurons potentiated cocaine-induced activity in male and female mice, consistent with studies involving DA neuron-specific ablation of D₂R in mice (Bello et al., 2011), and an RNAi-based approach

targeting D₂R in the rat VTA (de Jong et al., 2015). GABA_BR ablation also potentiated cocaine-induced activity, but the influence of this signaling pathway was restricted to males. Enhanced D₂R-dependent signaling in VTA DA neurons from female mice may compensate for the loss of GABA_BR signaling, explaining the weak influence of GABA_BR on cocaine-induced activity in females. Alternatively, there may be reduced GABAergic feedback to VTA DA neurons in females, rendering the loss of GABA_BR less effective (Zachry et al., 2021).

Behavioral sensitivity to morphine was unaffected by the loss of D₂R and/or GABA_BR in VTA DA neurons, consistent with previous reports (Steketee and Kalivas, 1991; Maldonado et al., 1997; Edwards et al., 2017). The lack of impact of D₂R ablation in VTA DA neurons is surprising given that opioids, like cocaine, increase VTA DA levels (Chefer et al., 2009). Cocaine and opioids differ, however,




Figure 4. Impact of GABA_BR and D₂R ablation on the motor-stimulatory effect of morphine. **A**, Total distance traveled during the 60-min period after injection of saline (left) or morphine (10 mg/kg, i.p., right) in female DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry ($n = 8$), AAV8-U6-gD2R-hSyn-NLSmCherry ($n = 8$), or AAV8-U6-gLacZ-hSyn-NLSmCherry control ($n = 8$). Two-way repeated measures ANOVA revealed a main effect of drug ($F_{(1,21)} = 185.3$, $p < 0.0001$), but no main effect of viral treatment ($F_{(2,21)} = 2.145$, $p = 0.1420$) or interaction between drug and viral treatment ($F_{(2,21)} = 1.183$, $p = 0.3260$); $+++p < 0.001$ (main effect of drug). **B**, Distance traveled for female subjects before and after morphine injection (denoted by arrow), evaluated in 5-min bins. Two-way repeated measures ANOVA revealed a main effect of bin ($F_{(3,057,67,26)} = 97.73$, $p < 0.0001$), but no effect of viral treatment ($F_{(2,22)} = 1.920$, $p = 0.1704$) or interaction between bin and viral treatment ($F_{(24,264)} = 0.5380$, $p = 0.9640$). **C**, Thigmotaxis index for female subjects following saline or morphine injection. Two-way repeated measures ANOVA revealed a main effect of drug ($F_{(1,21)} = 59.23$, $p < 0.0001$), but no significant difference with viral treatment ($F_{(2,21)} = 2.462$, $p = 0.1095$), and no interaction between drug and viral treatment ($F_{(2,21)} = 1.995$, $p = 0.1609$); $+++p < 0.001$ (main effect of drug). **D**, Total distance traveled during the 60-min period after injection of saline (left) or morphine (10 mg/kg, i.p., right) in male DATCre(+):Cas9GFP(+) mice treated with AAV8-U6-gGBR1-hSyn-NLSmCherry ($n = 7$), AAV8-U6-gD2R-hSyn-NLSmCherry ($n = 7$), or AAV8-U6-gLacZ-hSyn-NLSmCherry control ($n = 8$). Two-way repeated measures ANOVA revealed main effects of drug ($F_{(1,19)} = 176.1$, $p < 0.0001$), but no main effect of viral treatment ($F_{(2,19)} = 0.05804$, $p = 0.9438$) or interaction between drug and viral treatment ($F_{(2,19)} = 0.3193$, $p = 0.7305$); $+++p < 0.001$ (main effect of drug). **E**, Distance traveled for male subjects before and after morphine injection (denoted by arrow), evaluated in 5-min bins. Two-way repeated measures ANOVA revealed a main effect of bin ($F_{(2,264,43,02)} = 80.68$, $p < 0.0001$), but no main effect of viral treatment ($F_{(2,19)} = 0.1417$, $p = 0.8688$) or interaction between drug and viral treatment ($F_{(24,228)} = 0.9227$, $p = 0.5711$). **F**, Thigmotaxis index for male subjects following saline or morphine injection. Two-way repeated measures ANOVA revealed a main effect of drug ($F_{(1,19)} = 68.91$, $p < 0.0001$), but no significant difference with viral treatment ($F_{(2,19)} = 0.3907$, $p = 0.6819$), and no interaction between drug and viral treatment ($F_{(2,19)} = 1.195$, $p = 0.3244$); $+++p < 0.001$ (main effect of drug).

in their influence on VTA DA neurons. While cocaine hyperpolarizes VTA DA neurons in a D₂R-dependent manner (Beckstead et al., 2004), opioids increase VTA DA neuron firing by suppressing GABAergic input from local GABA neurons and/or RMTg GABA neurons (Johnson and North, 1992; Zhou et al., 2009; Jalabert et al., 2011), as well as disinhibiting glutamatergic input to VTA DA neurons (Yang et al., 2020). Thus, any inhibitory influence of somatodendritic D₂R activation triggered by the opioid-induced rise in VTA DA is likely offset by the excitatory influence of disinhibition.

Since morphine increases NAc DA levels (Spielewoy et al., 2000; Chefer et al., 2003; Vander Weele et al., 2014), and DA neurotransmission in the NAc drives GABA_BR-dependent feedback to VTA DA neurons (Edwards et al., 2017), it is also surprising that GABA_BR ablation does not impact behavioral sensitivity to morphine. This outcome

might reflect differences in the amplitude of the DA increase evoked by cocaine and morphine. Indeed, NAc DA levels in freely moving rats increased more in response to intravenous cocaine than morphine (Pontieri et al., 1995). The spatiotemporal pattern of DA increases in the NAc may also differ for cocaine and morphine. Consistent with this premise, cocaine elicited a more pronounced increase in DA in the NAc shell as compared with core (Aragona et al., 2008, 2009), whereas morphine evoked similar DA increases in NAc core and shell (Vander Weele et al., 2014). Moreover, the morphine-induced increase in NAc DA was relatively transient, a phenomenon potentially linked to a simultaneous increase in NAc GABA levels evoked by morphine (Vander Weele et al., 2014).

The differential apparent engagement of GABA_BR-dependent and D₂R-dependent signaling pathways by cocaine and morphine may also reflect differential molecular

target location. Morphine can act on opioid receptors in other brain regions to regulate motor activity, bypassing VTA DA neurons. For example, locomotor activity decreased initially following intra-NAc injection of morphine in rats (Costall et al., 1978). In addition, NAc lesions failed to eliminate morphine-induced motor activation, suggesting a potential role of other brain regions in this effect (Stevens et al., 1986). Also, intra-NAc infusion of morphine abolished the motor stimulatory effect of intra-NAc DA, showing that opioid and dopaminergic pathways in the NAc exert competing influence on locomotion (Layer et al., 1991).

The direct inhibitory influence of GABA_BR and D₂R activation on VTA DA neurons is mediated primarily by activation of GIRK channels (Brodie and Dunwiddie, 1990; Ackerman et al., 1993; Mercuri et al., 1997; Chen and Pan, 2000; Bunney et al., 2001; Beckstead et al., 2004; Cruz et al., 2004; Ford, 2014), though other effectors are modulated as well (Philippart and Khaliq, 2018; Su et al., 2019). Genetic ablation of *Girk2* globally or selectively in DA neurons correlated with increased motor-stimulatory effect of cocaine (Pravetoni and Wickman, 2008; Kotecki et al., 2015; McCall et al., 2017, 2019) and morphine (Kotecki et al., 2015). The contribution of GIRK channels to cocaine-induced activity was further localized to VTA DA neurons (McCall et al., 2019). Given that GIRK channels mediate the D₂R-dependent and GABA_BR-dependent inhibition of VTA DA neurons and that DA neuron-specific loss of GIRK channels enhances the motor-stimulatory effect of cocaine and morphine, the lack of impact of D₂R or GABA_BR ablation on morphine-induced activity was unexpected. It is possible that GABA_BR-dependent and D₂R-dependent signaling pathways are functionally redundant and that both need to be eliminated to see an influence on morphine-induced activity. GIRK channels in non-VTA DA neurons, perhaps in the adjacent substantia nigra pars compacta (Koyrakh et al., 2005), may also explain the impact of GIRK channel ablation on morphine.

Females are more susceptible to various facets of addiction (Fattore et al., 2008; Anker and Carroll, 2011; Becker and Koob, 2016), fueling interest in identifying relevant sex differences at molecular and cellular levels. Here, we found multiple sex differences related to inhibitory G-protein signaling in VTA DA neurons. Loss of GABA_BR, for example, had minimal impact on behavioral sensitivity to cocaine in females, but significantly enhanced the motor-stimulatory effect of cocaine in males. D₂R-dependent somatodendritic inhibitory currents were also larger in VTA DA neurons from females. This difference could reflect elevated D₂R expression and/or function in VTA DA neurons from females and is predicted to decrease cocaine sensitivity. Indeed, females exhibit elevated D₂R-dependent signaling at baseline as compared with males (Walker et al., 2006; Zachry et al., 2021).

Cocaine-induced activity was characterized by a more rapid and sharp peak in females as compared with males, which could be explained by a tighter regulation of synaptic DA by DAT in females compared with males (Zachry et al., 2021). D₂R ablation in VTA DA neurons in male mice yielded a temporal profile comparable to that seen in

female mice. While this suggests that D₂R-dependent signaling in VTA DA neurons is critical in tempering the early behavioral response to cocaine, the shift in the temporal profile observed in males is somewhat counterintuitive given that somatodendritic D₂R-dependent signaling is weaker in VTA DA neurons from males. This apparent discrepancy is perhaps explained by sex differences in the influence of presynaptic/terminal D₂R (which should also be eliminated by the CRISPR/Cas9 ablation approach) on DA dynamics in the NAc. Presynaptic D₂R-dependent signaling in VTA DA neurons in males may be stronger than that seen in females. Alternatively, while D₂R-dependent somatodendritic response amplitudes are larger in VTA DA neurons from female mice, VTA DA neurons in males may be more sensitive to D₂R-dependent inhibition.

While D₂R-dependent and GABA_BR-dependent signaling pathways in the VTA can suppress behavioral sensitivity to cocaine, psychostimulant exposure can weaken these inhibitory feedback pathways. For example, a transient decrease in D₂R influence on VTA DA neuron firing has been reported after repeated cocaine treatment or self-administration (Henry et al., 1989; Ackerman and White, 1990). This adaptation, and corresponding potentiation of cocaine-induced motor activity, was reproduced by repeated quinpirole treatment (Henry et al., 1998). Cocaine self-administration in rats also increased firing rate and burst activity of midbrain DA neurons, paralleled by decreased ability of quinpirole to inhibit DA neuron firing rate (Marinelli et al., 2003). Self-administration of amphetamine in rats reduced the ability of D₂R to suppress evoked DA release in the NAc, an effect mediated in part by an RGS2-dependent disruption of D_{2/3}R/Gα₁₂ functional coupling (Calipari et al., 2014; Sun et al., 2015). Repeated cocaine in male rats also decreased GABA_BR/G-protein coupling (Kushner and Unterwald, 2001). Finally, cocaine suppressed GABA_BR-dependent somatodendritic signaling in putative VTA DA neurons (Arora et al., 2011), and methamphetamine self-administration suppressed D₂R-dependent and GABA_BR-dependent somatodendritic responses in VTA DA neurons (Sharpe et al., 2014).

Notably, D₂R activation appears critical for many of these forms of plasticity, which likely contribute to the hyperexcitability of meso-accumbens DA pathway following psychostimulant exposure (Henry et al., 1998; Francis et al., 2019). The reciprocal relationship between psychostimulants and inhibitory G-protein-dependent feedback pathways in VTA DA neurons may be dependent on age and/or species, or methodological variables. Indeed, repeated methamphetamine injections correlated with enhanced D₂R-dependent hyperpolarization in young (8–10 d) rats (Amano et al., 2003). In mice, repeated methamphetamine injections suppressed somatodendritic GABA_BR-dependent (but not D₂R-dependent) signaling in VTA DA neurons, but only when methamphetamine was given in novel environment (Munoz et al., 2016).

Our work highlights innate signaling mechanisms that regulate behavioral sensitivity to cocaine. Knowledge of the molecular players and neuron populations regulating behavioral sensitivity to cocaine and other drugs of abuse

may help in identifying individuals at risk for developing addiction. Further investigation of mechanisms regulating the strength and sensitivity of inhibitory G-protein signaling pathways in VTA DA neurons may suggest opportunities for selective therapeutic interventions tailored to specific drugs of abuse.

References

- Ackerman JM, White FJ (1990) A10 somatodendritic dopamine autoreceptor sensitivity following withdrawal from repeated cocaine treatment. *Neurosci Lett* 117:181–187.
- Ackerman JM, Johansen PA, Clark D, White FJ (1993) Electrophysiological effects of putative autoreceptor-selective dopamine agonists on A10 dopamine neurons. *J Pharmacol Exp Ther* 265:963–970.
- Adell A, Artigas F (2004) The somatodendritic release of dopamine in the ventral tegmental area and its regulation by afferent transmitter systems. *Neurosci Biobehav Rev* 28:415–431.
- Adrover MF, Shin JH, Alvarez VA (2014) Glutamate and dopamine transmission from midbrain dopamine neurons share similar release properties but are differentially affected by cocaine. *J Neurosci* 34:3183–3192.
- Amano T, Matsubayashi H, Seki T, Sasa M, Sakai N (2003) Repeated administration of methamphetamine causes hypersensitivity of D2 receptor in rat ventral tegmental area. *Neurosci Lett* 347:89–92.
- Anker JJ, Carroll ME (2011) Females are more vulnerable to drug abuse than males: evidence from preclinical studies and the role of ovarian hormones. *Curr Top Behav Neurosci* 8:73–96.
- Aragona BJ, Cleaveland NA, Stuber GD, Day JJ, Carelli RM, Wightman RM (2008) Preferential enhancement of dopamine transmission within the nucleus accumbens shell by cocaine is attributable to a direct increase in phasic dopamine release events. *J Neurosci* 28:8821–8831.
- Aragona BJ, Day JJ, Roitman MF, Cleaveland NA, Wightman RM, Carelli RM (2009) Regional specificity in the real-time development of phasic dopamine transmission patterns during acquisition of a cue-cocaine association in rats. *Eur J Neurosci* 30:1889–1899.
- Arora D, Haluk DM, Kourrich S, Pravetoni M, Fernández-Alacid L, Nicolau JC, Luján R, Wickman K (2010) Altered neurotransmission in the mesolimbic reward system of Girk mice. *J Neurochem* 114:1487–1497.
- Arora D, Hearing M, Haluk DM, Mirkovic K, Fajardo-Serrano A, Wessendorf MW, Watanabe M, Luján R, Wickman K (2011) Acute cocaine exposure weakens GABA(B) receptor-dependent G-protein-gated inwardly rectifying K⁺ signaling in dopamine neurons of the ventral tegmental area. *J Neurosci* 31:12251–12257.
- Backes EN, Hemby SE (2008) Contribution of ventral tegmental GABA receptors to cocaine self-administration in rats. *Neurochem Res* 33:459–467.
- Beart PM, McDonald D, Gundlach AL (1979) Mesolimbic dopaminergic neurones and somatodendritic mechanisms. *Neurosci Lett* 15:165–170.
- Becker JB, Koob GF (2016) Sex differences in animal models: focus on addiction. *Pharmacol Rev* 68:242–263.
- Beckstead MJ, Grandy DK, Wickman K, Williams JT (2004) Vesicular dopamine release elicits an inhibitory postsynaptic current in mid-brain dopamine neurons. *Neuron* 42:939–946.
- Bello EP, Mateo Y, Gelman DM, Noain D, Shin JH, Low MJ, Alvarez VA, Lovinger DM, Rubinstein M (2011) Cocaine supersensitivity and enhanced motivation for reward in mice lacking dopamine D2 autoreceptors. *Nat Neurosci* 14:1033–1038.
- Bernosky-Smith KA, Qiu YY, Feja M, Lee YB, Loughlin B, Li JX, Bass CE (2018) Ventral tegmental area D2 receptor knockdown enhances choice impulsivity in a delay-discounting task in rats. *Behav Brain Res* 341:129–134.
- Boender AJ, de Jong JW, Boekhoudt L, Luijendijk MC, van der Plasse G, Adan RA (2014) Combined use of the canine adenovirus-2 and DREADD-technology to activate specific neural pathways in vivo. *PLoS One* 9:e95392.
- Bradberry CW, Roth RH (1989) Cocaine increases extracellular dopamine in rat nucleus accumbens and ventral tegmental area as shown by in vivo microdialysis. *Neurosci Lett* 103:97–102.
- Brebner K, Phelan R, Roberts DC (2000) Intra-VTA baclofen attenuates cocaine self-administration on a progressive ratio schedule of reinforcement. *Pharmacol Biochem Behav* 66:857–862.
- Brodie MS, Dunwiddie TV (1990) Cocaine effects in the ventral tegmental area: evidence for an indirect dopaminergic mechanism of action. *Naunyn Schmiedeberg Arch Pharmacol* 342:660–665.
- Bunney EB, Appel SB, Brodie MS (2001) Electrophysiological effects of cocaethylene, cocaine, and ethanol on dopaminergic neurons of the ventral tegmental area. *J Pharmacol Exp Ther* 297:696–703.
- Calipari ES, Sun H, Eldeeb K, Luessen DJ, Feng X, Howlett AC, Jones SR, Chen R (2014) Amphetamine self-administration attenuates dopamine D2 autoreceptor function. *Neuropsychopharmacology* 39:1833–1842.
- Chefer VI, Kieffer BL, Shippenberg TS (2003) Basal and morphine-evoked dopaminergic neurotransmission in the nucleus accumbens of MOR- and DOR-knockout mice. *Eur J Neurosci* 18:1915–1922.
- Chefer VI, Denoroy L, Zapata A, Shippenberg TS (2009) Mu opioid receptor modulation of somatodendritic dopamine overflow: GABAergic and glutamatergic mechanisms. *Eur J Neurosci* 30:272–278.
- Chen NH, Reith ME (1994) Autoregulation and monoamine interactions in the ventral tegmental area in the absence and presence of cocaine: a microdialysis study in freely moving rats. *J Pharmacol Exp Ther* 271:1597–1610.
- Chen NN, Pan WH (2000) Regulatory effects of D2 receptors in the ventral tegmental area on the mesocorticolimbic dopaminergic pathway. *J Neurochem* 74:2576–2582.
- Chen R, McIntosh S, Hemby SE, Sun H, Sexton T, Martin TJ, Childers SR (2018) High and low doses of cocaine intake are differentially regulated by dopamine D2 receptors in the ventral tegmental area and the nucleus accumbens. *Neurosci Lett* 671:133–139.
- Chen SH, Haam J, Walker M, Scappini E, Naughton J, Martin NP (2019) Production of viral constructs for neuroanatomy, calcium imaging, and optogenetics. *Curr Protoc Neurosci* 87:e66.
- Chen SY, Burger RI, Reith ME (1996) Extracellular dopamine in the rat ventral tegmental area and nucleus accumbens following ventral tegmental infusion of cocaine. *Brain Res* 729:294–296.
- Costall B, Fortune DH, Naylor RJ (1978) The induction of catalepsy and hyperactivity by morphine administered directly into the nucleus accumbens of rats. *Eur J Pharmacol* 49:49–64.
- Cruz HG, Ivanova T, Lunn M-L, Stoffel M, Slesinger PA, Lüscher C (2004) Bi-directional effects of GABA(B) receptor agonists on the mesolimbic dopamine system. *Nat Neurosci* 7:153–159.
- de Jong JW, Roelofs TJ, Mol FM, Hillen AE, Meijboom KE, Luijendijk MC, van der Eerden HA, Garner KM, Vanderschuren LJ, Adan RA (2015) Reducing ventral tegmental dopamine D2 receptor expression selectively boosts incentive motivation. *Neuropsychopharmacology* 40:2085–2095.
- Delfs JM, Schreiber L, Kelley AE (1990) Microinjection of cocaine into the nucleus accumbens elicits locomotor activation in the rat. *J Neurosci* 10:303–310.
- Di Chiara G, Imperato A (1988) Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci USA* 85:5274–5278.
- Edwards NJ, Tejada HA, Pignatelli M, Zhang S, McDevitt RA, Wu J, Bass CE, Bettler B, Morales M, Bonci A (2017) Circuit specificity in the inhibitory architecture of the VTA regulates cocaine-induced behavior. *Nat Neurosci* 20:438–448.
- Einhorn LC, Johansen PA, White FJ (1988) Electrophysiological effects of cocaine in the mesoaccumbens dopamine system: studies in the ventral tegmental area. *J Neurosci* 8:100–112.

- Fattore L, Altea S, Fratta W (2008) Sex differences in drug addiction: a review of animal and human studies. *Womens Health (Lond)* 4:51–65.
- Ford CP (2014) The role of D2-autoreceptors in regulating dopamine neuron activity and transmission. *Neuroscience* 282:13–22.
- Francis TC, Gantz SC, Moussawi K, Bonci A (2019) Synaptic and intrinsic plasticity in the ventral tegmental area after chronic cocaine. *Curr Opin Neurobiol* 54:66–72.
- Groves PM, Wilson CJ, Young SJ, Rebec GV (1975) Self-inhibition by dopaminergic neurons. *Science* 190:522–528.
- Guo S, Chen S, Zhang Q, Wang Y, Xu K, Zheng X (2014) Optogenetic activation of the excitatory neurons expressing CaMKII α in the ventral tegmental area upregulates the locomotor activity of free behaving rats. *Biomed Res Int* 2014:687469.
- Henry DJ, Greene MA, White FJ (1989) Electrophysiological effects of cocaine in the mesoaccumbens dopamine system: repeated administration. *J Pharmacol Exp Ther* 251:833–839.
- Henry DJ, Hu XT, White FJ (1998) Adaptations in the mesoaccumbens dopamine system resulting from repeated administration of dopamine D1 and D2 receptor-selective agonists: relevance to cocaine sensitization. *Psychopharmacology (Berl)* 140:233–242.
- Holroyd KB, Adrover MF, Fuino RL, Bock R, Kaplan AR, Gremel CM, Rubinstein M, Alvarez VA (2015) Loss of feedback inhibition via D2 autoreceptors enhances acquisition of cocaine taking and reactivity to drug-paired cues. *Neuropsychopharmacology* 40:1495–1509.
- Hyman SE, Malenka RC, Nestler EJ (2006) Neural mechanisms of addiction: the role of reward-related learning and memory. *Annu Rev Neurosci* 29:565–598.
- Iravani MM, Muscat R, Kruk ZL (1996) Comparison of somatodendritic and axon terminal dopamine release in the ventral tegmental area and the nucleus accumbens. *Neuroscience* 70:1025–1037.
- Jalabert M, Bourdy R, Courtin J, Veinante P, Manzoni OJ, Barrot M, Georges F (2011) Neuronal circuits underlying acute morphine action on dopamine neurons. *Proc Natl Acad Sci USA* 108:16446–16450.
- Jhou TC, Fields HL, Baxter MG, Saper CB, Holland PC (2009) The rostromedial tegmental nucleus (RMTg), a GABAergic afferent to midbrain dopamine neurons, encodes aversive stimuli and inhibits motor responses. *Neuron* 61:786–800.
- Johnson SW, North RA (1992) Opioids excite dopamine neurons by hyperpolarization of local interneurons. *J Neurosci* 12:483–488.
- Juarez B, Han MH (2016) Diversity of dopaminergic neural circuits in response to drug exposure. *Neuropsychopharmacology* 41:2424–2446.
- Kalivas PW, Duffy P (1993) Time course of extracellular dopamine and behavioral sensitization to cocaine. I. Dopamine axon terminals. *J Neurosci* 13:266–275.
- Kalivas PW, Duffy P, Eberhardt H (1990) Modulation of A10 dopamine neurons by gamma-aminobutyric acid agonists. *J Pharmacol Exp Ther* 253:858–866.
- Kim KM, Baratta MV, Yang A, Lee D, Boyden ES, Fiorillo CD (2012) Optogenetic mimicry of the transient activation of dopamine neurons by natural reward is sufficient for operant reinforcement. *PLoS One* 7:e33612.
- Kita JM, Kile BM, Parker LE, Wightman RM (2009) In vivo measurement of somatodendritic release of dopamine in the ventral tegmental area. *Synapse* 63:951–960.
- Kotecki L, Hearing M, McCall NM, Marron Fernandez de Velasco E, Pravetoni M, Arora D, Victoria NC, Munoz MB, Xia Z, Slesinger PA, Weaver CD, Wickman K (2015) GIRK channels modulate opioid-induced motor activity in a cell type- and subunit-dependent manner. *J Neurosci* 35:7131–7142.
- Koulchitsky S, Delaïresse C, Beeken T, Monteforte A, Dethier J, Quertermont E, Findeisen R, Bullinger E, Seutin V (2016) Activation of D2 autoreceptors alters cocaine-induced locomotion and slows down local field oscillations in the rat ventral tegmental area. *Neuropharmacology* 108:120–127.
- Koyrakh L, Luján R, Colón J, Karschin C, Kurachi Y, Karschin A, Wickman K (2005) Molecular and cellular diversity of neuronal G-protein-gated potassium channels. *J Neurosci* 25:11468–11478.
- Kushner SA, Unterwald EM (2001) Chronic cocaine administration decreases the functional coupling of GABA(B) receptors in the rat ventral tegmental area as measured by baclofen-stimulated 35S-GTPgammaS binding. *Life Sci* 69:1093–1102.
- Labouèbe G, Lomazzi M, Cruz HG, Creton C, Luján R, Li M, Yanagawa Y, Obata K, Watanabe M, Wickman K, Boyer SB, Slesinger PA, Lüscher C (2007) RGS2 modulates coupling between GABAB receptors and GIRK channels in dopamine neurons of the ventral tegmental area. *Nat Neurosci* 10:1559–1568.
- Lammel S, Lim BK, Malenka RC (2014) Reward and aversion in a heterogeneous midbrain dopamine system. *Neuropharmacology* 76:351–359.
- Layer RT, Uretsky NJ, Wallace LJ (1991) Effects of morphine in the nucleus accumbens on stimulant-induced locomotion. *Pharmacol Biochem Behav* 40:21–26.
- Leite-Morris KA, Fukudome EY, Kaplan GB (2002) Opiate-induced motor stimulation is regulated by gamma-aminobutyric acid type B receptors found in the ventral tegmental area in mice. *Neurosci Lett* 317:119–122.
- Leite-Morris KA, Fukudome EY, Shoeb MH, Kaplan GB (2004) GABA (B) receptor activation in the ventral tegmental area inhibits the acquisition and expression of opiate-induced motor sensitization. *J Pharmacol Exp Ther* 308:667–678.
- Lüscher C (2016) The emergence of a circuit model for addiction. *Annu Rev Neurosci* 39:257–276.
- Lüscher C, Ungless MA (2006) The mechanistic classification of addictive drugs. *PLoS Med* 3:e437. [
- Maldonado R, Saiardi A, Valverde O, Samad TA, Roques BP, Borrelli E (1997) Absence of opiate rewarding effects in mice lacking dopamine D2 receptors. *Nature* 388:586–589.
- Marinelli M, Cooper DC, Baker LK, White FJ (2003) Impulse activity of midbrain dopamine neurons modulates drug-seeking behavior. *Psychopharmacology (Berl)* 168:84–98.
- Martin TA, Smith HR, Luessen DJ, Chen R, Porrino LJ (2020) Functional brain activity is globally elevated by dopamine D2 receptor knockdown in the ventral tegmental area. *Brain Res* 1727:146552.
- McCall NM, Kotecki L, Dominguez-Lopez S, Marron Fernandez de Velasco E, Carlblom N, Sharpe AL, Beckstead MJ, Wickman K (2017) Selective ablation of GIRK channels in dopamine neurons alters behavioral effects of cocaine in mice. *Neuropsychopharmacology* 42:707–715.
- McCall NM, Marron Fernandez de Velasco E, Wickman K (2019) GIRK channel activity in dopamine neurons of the ventral tegmental area bidirectionally regulates behavioral sensitivity to cocaine. *J Neurosci* 39:3600–3610.
- Menegas W, Bergan JF, Ogawa SK, Isogai Y, Umadevi Venkataraju K, Osten P, Uchida N, Watabe-Uchida M (2015) Dopamine neurons projecting to the posterior striatum form an anatomically distinct subclass. *Elife* 4:e10032.
- Mercuri NB, Saiardi A, Bonci A, Picetti R, Calabresi P, Bernardi G, Borrelli E (1997) Loss of autoreceptor function in dopaminergic neurons from dopamine D2 receptor deficient mice. *Neuroscience* 79:323–327.
- Mickley GA, Mulvihill MA, Postler MA (1990) Brain mu and delta opioid receptors mediate different locomotor hyperactivity responses of the C57BL/6J mouse. *Psychopharmacology (Berl)* 101:332–337.
- Munoz MB, Padgett CL, Rifkin R, Terunuma M, Wickman K, Contet C, Moss SJ, Slesinger PA (2016) A role for the GIRK3 subunit in methamphetamine-induced attenuation of GABAB receptor-activated GIRK currents in VTA dopamine neurons. *J Neurosci* 36:3106–3114.
- Narayanan S, Wallace L, Uretsky N (1996) Spontaneous and drug-stimulated locomotor activity after the administration of pertussis toxin into the ventral tegmental area. *J Psychiatry Neurosci* 21:172–180.

- Philippart F, Khaliq ZM (2018) Gi/o protein-coupled receptors in dopamine neurons inhibit the sodium leak channel NALCN. *Elife* 7:e40984.
- Pignatelli M, Bonci A (2018) Spiraling connectivity of NAc-VTA circuitry. *Neuron* 97:261–262.
- Platt RJ, Chen S, Zhou Y, Yim MJ, Swiech L, Kempton HR, Dahlman JE, Parnas O, Eisenhaure TM, Jovanovic M, Graham DB, Jhunjhunwala S, Heidenreich M, Xavier RJ, Langer R, Anderson DG, Hacohen N, Regev A, Feng G, Sharp PA, et al. (2014) CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell* 159:440–455.
- Pontieri FE, Tanda G, Di Chiara G (1995) Intravenous cocaine, morphine, and amphetamine preferentially increase extracellular dopamine in the “shell” as compared with the “core” of the rat nucleus accumbens. *Proc Natl Acad Sci USA* 92:12304–12308.
- Pravetoni M, Wickman K (2008) Behavioral characterization of mice lacking GIRK/Kir3 channel subunits. *Genes Brain Behav* 7:523–531.
- Rahman S, McBride WJ (2000) Feedback control of mesolimbic somatodendritic dopamine release in rat brain. *J Neurochem* 74:684–692.
- Rahman S, McBride WJ (2001) D1-D2 dopamine receptor interaction within the nucleus accumbens mediates long-loop negative feedback to the ventral tegmental area (VTA). *J Neurochem* 77:1248–1255.
- Runegaard AH, Sørensen AT, Fitzpatrick CM, Jørgensen SH, Petersen AV, Hansen NW, Weikop P, Andreassen JT, Mikkelsen JD, Perrier JF, Woldbye D, Rickhag M, Wortwein G, Gether U (2018) Locomotor- and reward-enhancing effects of cocaine are differentially regulated by chemogenetic stimulation of Gi-signaling in dopaminergic neurons. *eNeuro* 5:ENEURO.0345-17.2018.
- Schultz W (2016) Dopamine reward prediction-error signalling: a two-component response. *Nat Rev Neurosci* 17:183–195.
- Sharpe AL, Varela E, Bettinger L, Beckstead MJ (2014) Methamphetamine self-administration in mice decreases GIRK channel-mediated currents in midbrain dopamine neurons. *Int J Neuropsychopharmacol* 18:pyu073–pyu073.
- Spielewoy C, Gonon F, Roubert C, Fauchey V, Jaber M, Caron MG, Roques BP, Hamon M, Betancur C, Maldonado R, Giros B (2000) Increased rewarding properties of morphine in dopamine-transporter knockout mice. *Eur J Neurosci* 12:1827–1837.
- Steketee JD, Kalivas PW (1990) Sensitization to cocaine produced by injection of pertussis toxin into the A10 dopamine region. *NIDA Res Monogr* 105:545–546.
- Steketee JD, Kalivas PW (1991) Sensitization to psychostimulants and stress after injection of pertussis toxin into the A10 dopamine region. *J Pharmacol Exp Ther* 259:916–924.
- Stevens KE, Mickley GA, McDermott LJ (1986) Brain areas involved in production of morphine-induced locomotor hyperactivity of the C57B1/6J mouse. *Pharmacol Biochem Behav* 24:1739–1747.
- Su M, Li L, Wang J, Sun H, Zhang L, Zhao C, Xie Y, Gamper N, Du X, Zhang H (2019) Kv7.4 channel contribute to projection-specific auto-inhibition of dopamine neurons in the ventral tegmental area. *Front Cell Neurosci* 13:557.
- Sun H, Calipari ES, Beveridge TJ, Jones SR, Chen R (2015) The brain gene expression profile of dopamine D2/D3 receptors and associated signaling proteins following amphetamine self-administration. *Neuroscience* 307:253–261.
- Tanabe LM, Suto N, Creekmore E, Steinmiller CL, Vezina P (2004) Blockade of D2 dopamine receptors in the VTA induces a long-lasting enhancement of the locomotor activating effects of amphetamine. *Behav Pharmacol* 15:387–395.
- Tye KM, Mirzabekov JJ, Warden MR, Ferenczi EA, Tsai HC, Finkelstein J, Kim SY, Adhikari A, Thompson KR, Andalman AS, Gunaydin LA, Witten IB, Deisseroth K (2013) Dopamine neurons modulate neural encoding and expression of depression-related behaviour. *Nature* 493:537–541.
- van Rossum J, van der SJ, Hurkmans JA (1962) Mechanism of action of cocaine and amphetamine in the brain. *Experientia* 18:229–231.
- Vander Weele CM, Porter-Stransky KA, Mabrouk OS, Lovic V, Singer BF, Kennedy RT, Aragona BJ (2014) Rapid dopamine transmission within the nucleus accumbens: dramatic difference between morphine and oxycodone delivery. *Eur J Neurosci* 40:3041–3054.
- Volkow ND, Morales M (2015) The brain on drugs: from reward to addiction. *Cell* 162:712–725.
- Walker QD, Ray R, Kuhn CM (2006) Sex differences in neurochemical effects of dopaminergic drugs in rat striatum. *Neuropsychopharmacology* 31:1193–1202.
- Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N (2012) Whole-brain mapping of direct inputs to midbrain dopamine neurons. *Neuron* 74:858–873.
- Xi ZX, Stein EA (1999) Baclofen inhibits heroin self-administration behavior and mesolimbic dopamine release. *J Pharmacol Exp Ther* 290:1369–1374.
- Xue Y, Steketee JD, Rebec GV, Sun W (2011) Activation of D2-like receptors in rat ventral tegmental area inhibits cocaine-reinstated drug-seeking behavior. *Eur J Neurosci* 33:1291–1298.
- Yang L, Chen M, Ma Q, Sheng H, Cui D, Shao D, Lai B, Zheng P (2020) Morphine selectively disinhibits glutamatergic input from mPFC onto dopamine neurons of VTA, inducing reward. *Neuropharmacology* 176:108217.
- Zachry JE, Nolan SO, Brady LJ, Kelly SJ, Siciliano CA, Calipari ES (2021) Sex differences in dopamine release regulation in the striatum. *Neuropsychopharmacology* 46:491–499.