

Disorders of the Nervous System

Highlighted Research Paper: Chemogenetic Suppression of Medial Prefrontal-Dorsal Hippocampal Interactions Prevents Estrogenic Enhancement of Memory Consolidation in Female Mice by, Jennifer J. Tuscher, Lisa R. Taxier, Jayson C. Schalk, Jacqueline M. Haertel, and Karyn M. Frick.

Concurrent Medial Prefrontal Cortex and Dorsal Hippocampal Activity Is Required for Estradiol-Mediated Effects on Object Memory and Spatial Memory Consolidation

Rosalind S.E. Carney, DPhil

Steroid hormones such as estrogens and androgens affect the nervous system during *in utero* development and throughout adolescence, adulthood, and aging (Choleris et al., 2018). Steroid hormones have been associated with a prevalence in either men or women of certain neurologic disorders. For instance, Alzheimer's disease and schizophrenia affect more women than men and vice versa, respectively (Irvine et al., 2012; Jackson et al., 2013). Studies have also shown that 17 β -estradiol (E₂), the predominant and most potent form of estrogen, can alter hippocampal neuroplasticity and therefore affect learning and memory. Exogenous E₂ administration into the dorsal hippocampus (DH) of young female rodents enhances memory consolidation as shown by object recognition (OR) and object placement (OP; also known as object location) tasks (Tuscher et al., 2015). OR and OP are two object-based one-trial learning tasks that capitalize on rodents' innate interest in novelty and relate to the "what" and "where" aspects of episodic memory consolidation and spatial memory, respectively (Frick et al., 2018a). DH E₂ infusion immediately after the OR/OP training period restricts the effects of E₂ to the time frame of memory consolidation (Tuscher et al., 2015). Post-training DH E₂ infusion leads to an increase in spine density of CA1 pyramidal neurons during the time frame in which memory consolidation occurs (Gould et al., 1990; Woolley et al., 1990; Woolley and McEwen, 1993; MacLusky et al., 2005; Phan et al., 2012). Increased dendritic spine density in response to post-training E₂ DH infusion was also observed in the medial prefrontal cortex (mPFC; Tuscher

et al., 2016), a structure that expresses estrogen receptors and is also involved in memory consolidation. This observation prompted Tuscher and colleagues to examine in their *eNeuro* publication the effects of local infusion of E₂ into the mPFC and the requirement for concurrent activity of the mPFC in E₂-mediated memory consolidation in the DH.

Young female mice were ovariectomized (OVX) to eliminate the ovarian source of E₂ that fluctuates during the estrous cycle, the rodent version of the menstrual cycle (Tuscher et al., 2015). Figure 1 shows a schematic illustration of the experimental approach. For OR training, a mouse was placed in a white square arena in which the upper left and right corners contained identical objects, one in each corner (Fig. 1A). The time spent exploring each object was recorded using tracking software, and mice remained in the arena until they had accumulated 30 s exploring the objects. If the mice remembered the identity of the training objects during testing 48 h later, then they should spend more time than chance (>15 s) exploring a novel object that had replaced one of the original training objects in the upper corners of the arena (Fig. 1A). The OP assessment differed from that of OR in two aspects: (1) OP testing occurred 24 h after training; and (2) whereas one of the training objects encountered during training remained in its original location during testing, the second training object had been moved to a lower corner of the arena (Fig. 1B). Spending more time than chance with the moved object indicated that spatial memory was intact.

OVX mice were implanted with bilateral cannulae targeted to the mPFC one week before training in the OR and OP tasks, which were performed in separate groups of mice. Immediately after training, mice received mPFC infusions of either cyclodextrin-encapsulated E₂ dissolved in sterile saline or the vehicle control, 2-hydroxypropyl- β -cyclodextrin (HBC) dissolved to the same concentration

Received July 12, 2019; accepted July 12, 2019; First published August 20, 2019.

<https://doi.org/10.1523/ENEURO.0271-19.2019>

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.




Figure 1. Schematic illustration of OR and OP behavioral assessments. **A**, Treatment administered after training reveals treatment effect on memory consolidation of a novel object versus a familiar object. **B**, Treatment administered after training reveals treatment effect on memory consolidation of the placement of a previously encountered object in a new location versus original placement of another previously encountered object (Adapted from Figure 1 in [Tuscher et al., 2019](#)).

of cyclodextrin as the E_2 -containing solution. In both the OR and OP behavioral assessments, E_2 -infused mice spent more time exploring the novel or moved objects than would be expected by chance alone and significantly more time compared to vehicle-infused mice ([Fig. 2](#)). These observations showed that local effects of E_2 within the mPFC replicate the enhancement of memory consolidation and spatial memory that resulted from local infusion of E_2 within the DH.

Although a causal link between E_2 -mediated spinogenesis and memory consolidation has not yet been established ([Sheppard et al., 2019](#)), signaling mechanisms are common to both processes ([Frick et al., 2018a](#)). Therefore, [Tuscher](#) and colleagues examined local E_2 -mediated effects on spinogenesis in the mPFC. Two weeks after the completion of the OR and OP behavioral assessments, mice received an infusion of E_2 or vehicle control into the mPFC; 2 h later the brains of the mice were harvested and processed for Golgi staining. Dendritic spine analysis was performed in regions in which increased E_2 -mediated spinogenesis was previously observed, namely, Layer II/III of the mPFC and CA1 of the DH. Spine density was increased apically but not basally when E_2 was locally infused into the mPFC, but spine density in the DH was

unchanged. It is not known why local infusion of E_2 did not affect spinogenesis in the DH; perhaps the time frame was insufficient to detect any alterations or E_2 infusion directly into the DH has more hierarchical, rather than reciprocal, effects on another brain region.

The observations described thus far provided evidence that E_2 acting locally in the mPFC enhances memory consolidation and increases local spinogenesis. Next, the authors sought to determine if concurrent activation of the mPFC is required for enhanced E_2 -mediated, DH-dependent memory consolidation. To do so, they employed a designer receptors exclusively activated by designer drugs (DREADD) approach to suppress activity in the mPFC while simultaneously infusing E_2 into the DH. Immediately after ovariectomy, viral DREADD constructs or saline (sham condition) were infused bilaterally into the mPFC, and cannulae were also placed bilaterally in the DH for future infusions. The AAV-CaMKII α -eGFP construct induced expression of green fluorescent protein (GFP) in excitatory neurons in the mPFC three weeks later, at the time when behavioral assessments were initiated. At the time of the surgical procedure, another group of OVX mice received bilateral mPFC injections of an AAV-CaMKII α -HA-hM4Di-IRES-mCitrine viral construct. Binding of the




Figure 2. Infusion of E_2 into the mPFC immediately after training enhances memory consolidation. Mice infused with E_2 directly into the mPFC spent significantly more time than chance (dashed line at 15 s) with the novel object (**A**) when tested 48 h after training or with the moved object (**B**) 24 h after training. Mice infused with vehicle (Veh) into the mPFC did not spend more time than chance with the novel (**A**) or moved (**B**) objects. These data suggest that E_2 can improve the consolidation of object memories by acting directly within the mPFC. Bars represent the mean \pm SEM; * p < 0.05 relative to chance and the vehicle group (Adapted from Figure 1 in [Tuscher et al., 2019](#)).




Figure 3. Chemogenetic suppression of the mPFC immediately after training prevents the memory enhancement induced by DH E_2 infusion. Sham, GFP, or hM4Di control groups receiving CNO + Veh did not spend significantly more time than chance (15 s) with the novel object (**A**) when tested 48 h after training or with the moved object (**B**) 24 h after training. In contrast, sham or GFP mice receiving CNO + E_2 immediately after training did spend significantly more time than chance with the novel (**A**) and moved (**B**) objects, displaying intact OR and OP memory. However, hM4Di mice treated with CNO+ E_2 immediately after training did not demonstrate intact memory, suggesting that DREADD-mediated suppression of the mPFC blocks the beneficial mnemonic effects of DH-infused E_2 . Bars represent the mean \pm SEM; * p < 0.05 relative to chance, all Veh-infused groups, and the hM4Di- E_2 group; + p < 0.05 relative to chance and the sham + Veh-infused and hM4Di- E_2 groups (Adapted from Figure 4 in [Tuscher et al., 2019](#)).

clozapine-N-oxide (CNO) ligand to the hM4Di DREADD suppresses the activity of the excitatory CaMKII α -expressing cells, thereby inhibiting the mPFC. After training in the OR or OP tasks, mice received DH infusion of E_2 or vehicle control at the same time that CNO was administered systemically to all experimental groups, at a concentration of CNO that does not impair memory consolidation on its own ([Tuscher et al., 2018](#)). In both the OR and OP tasks, memory consolidation was enhanced in mice in which E_2 was infused directly into the DH but in which mPFC activity was not suppressed (sham + E_2 , GFP + E_2 groups; [Fig. 3](#)) compared to experimental groups that received vehicle control (sham + HBC, GFP + HBC, hM4Di + HBC; [Fig. 3](#)). Suppression of excitatory signaling in the mPFC blocked the E_2 -mediated DH-dependent enhancement of memory consolidation (hM4Di + HBC; [Fig. 3](#)) showing, for the first time, that mPFC activity is required for memory consolidation when E_2 is locally infused into the DH.

The *eNeuro* publication by Tuscher and colleagues highlights the complexity of E_2 -mediated memory consolidation processes and raises several new avenues of research. The finding that local E_2 infusion into the mPFC affects a cognitive process that is heavily associated with the hippocampus underscores the importance of research into other brain structures involved in cognition that express steroid receptors. The mPFC and DH interaction could be regulated by direct or indirect neural circuitry. It will also be interesting to determine if local effects of E_2 within the mPFC involve classic, genomic signaling via estrogen receptors ER α and ER β or the rapid signaling that results from extracellular ligand binding to the G protein-coupled receptor GPER. Coupling the use of estrogen agonists and antagonists in the DH with chemogenetic suppression of other brain regions will help

resolve the interactions between signaling mechanisms and neural circuitry. Such receptor manipulation could also reveal the relevance of local estradiol production within the hippocampus that is unaffected by ovariectomy.

Tuscher and colleagues' findings in this *eNeuro* publication may also help resolve discrepancies in the literature regarding the effects of E_2 on cognition by emphasizing the need to examine other brain regions involved in memory consolidation. But more complexities exist, such as the effects of steroid signaling in young versus older or female versus male rodent models ([Frick et al., 2018b](#)). It was expected that hormone replacement therapy in postmenopausal women would have a protective effect against cognitive dysfunction. However, some of these studies used conjugated equine estrogens, rather than E_2 , which may explain the lack of beneficial effects ([Luine, 2014](#)). Moreover, it has recently been shown in a mouse model of Alzheimer's disease that E_2 fluctuations during the ovarian cycle have adverse effects on cognition ([Broestl et al., 2018](#)). Therefore, a plethora of age, sex, and brain region considerations may be needed to fully elucidate new targets for drug development that address steroid-related effects on cognitive functions, such as memory consolidation.

References

- Broestl L, Worden K, Moreno AJ, Davis EJ, Wang D, Garay B, Singh T, Verret L, Palop JJ, Dubal DB (2018) Ovarian cycle stages modulate Alzheimer-related cognitive and brain network alterations in female mice. *eNeuro* 5: ENEURO.0132-17.2018.
- Choleris E, Galea LAM, Sohrabji F, Frick KM (2018) Sex differences in the brain: implications for behavioral and biomedical research. *Neurosci Biobehav Rev* 85:126–145.

- Frick KM, Tuscher JJ, Koss WA, Kim J, Taxier LR (2018a) Estrogenic regulation of memory consolidation: a look beyond the hippocampus, ovaries, and females. *Physiol Behav* 187:57–66.
- Frick KM, Kim J, Koss WA (2018b) Estradiol and hippocampal memory in female and male rodents. *Curr Opin Behav Sci* 23:65–74.
- Gould E, Woolley CS, Frankfurt M, McEwen BS (1990) Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *J Neurosci* 10:1286–1291.
- Irvine K, Laws KR, Gale TM, Kondel TK (2012) Greater cognitive deterioration in women than men with Alzheimer's disease: a meta analysis. *J Clin Exp Neuropsychol* 34:989–998.
- Jackson D, Kirkbride J, Croudace T, Morgan C, Boydell J, Errazuriz A, Murray RM, Jones PB (2013) Meta-analytic approaches to determine gender differences in the age-incidence characteristics of schizophrenia and related psychoses. *Int J Methods Psychiatr Res* 22:36–45.
- Luine VN (2014) Estradiol and cognitive function: past, present and future. *Horm Behav* 66:602–618.
- MacLusky NJ, Luine VN, Hajszan T, Leranth C (2005) The 17 α and 17 β isomers of estradiol both induce rapid spine synapse formation in the CA1 hippocampal subfield of ovariectomized female rats. *Endocrinology* 146:287–293.
- Phan A, Gabor CS, Favaro KJ, Kaschack S, Armstrong JN, MacLusky NJ, Choleris E (2012) Low doses of 17 β -estradiol rapidly improve learning and increase hippocampal dendritic spines. *Neuropsychopharmacology* 37:2299–2309.
- Sheppard PAS, Choleris E, Galea L (2019) Structural plasticity of the hippocampus in response to estrogens in female rodents. *Mol Brain* 12:22.
- Tuscher JJ, Fortress AM, Kim J, Frick KM (2015) Regulation of object recognition and object placement by ovarian sex steroid hormones. *Behav Brain Res* 285:140–157.
- Tuscher JJ, Luine V, Frankfurt M, Frick KM (2016) Estradiol-mediated spine changes in the dorsal hippocampus and medial prefrontal cortex of ovariectomized female mice depend on ERK and mTOR activation in the dorsal hippocampus. *J Neurosci* 36:1483–1489.
- Tuscher JJ, Taxier LR, Fortress AM, Frick KM (2018) Chemogenetic inactivation of the dorsal hippocampus and medial prefrontal cortex, individually and concurrently, impairs object recognition and spatial memory consolidation in female mice. *Neurobiol Learn Mem* 156:103–116.
- Tuscher JJ, Taxier LR, Schalk JC, Haertel JM, Frick KM (2019) Chemogenetic suppression of medial prefrontal-dorsal hippocampal interactions prevents estrogenic enhancement of memory consolidation in female mice. *eNeuro* 6: ENEURO.0451-18.2019
- Woolley CS, McEwen BS (1993) Roles of estradiol and progesterone in regulation of hippocampal dendritic spine density during the estrous cycle in the rat. *J Comp Neurol* 336:293–306.
- Woolley CS, Gould E, Frankfurt M, McEwen BS (1990) Naturally occurring fluctuation in dendritic spine density on adult hippocampal pyramidal neurons. *J Neurosci* 10:4035–4039.