

Research Article: New Research | Disorders of the Nervous System

Altered cortico-brainstem connectivity during spontaneous fluctuations in pain intensity in painful trigeminal neuropathy.

<https://doi.org/10.1523/ENEURO.0522-23.2024>

Received: 10 December 2023

Revised: 2 June 2024

Accepted: 6 June 2024

Copyright © 2024 Meylakh et al.

This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

This Early Release article has been peer reviewed and accepted, but has not been through the composition and copyediting processes. The final version may differ slightly in style or formatting and will contain links to any extended data.

Alerts: Sign up at www.eneuro.org/alerts to receive customized email alerts when the fully formatted version of this article is published.

1. Manuscript Title: Altered cortico-brainstem connectivity during spontaneous fluctuations in pain intensity in painful trigeminal neuropathy.

2. Abbreviated title: Fluctuating chronic neuropathic pain

3. Authors and affiliations: N. Meylakh¹, L.S. Crawford¹, E.P. Mills¹, V.G. Macefield², E.R. Vickers¹, P.M. Macey³, K.A. Keay¹, and L.A. Henderson¹.

¹ School of Medical Sciences (Neuroscience), Brain and Mind Centre, University of Sydney, Australia 2050; ²Department of Neuroscience, Monash University, Victoria, Australia 3800; ³UCLA School of Nursing and Brain Research Institute, University of California, Los Angeles, California, United States 90095.

4. Author Contributions: NM and LSC analyzed data and wrote the paper; EPM performed research; VGM wrote the paper; ERV and PMM contributed unpublished analytic tools; KAK and LAH designed research and wrote the paper.

5. Corresponding Author: Luke A Henderson, School of Medical Sciences (Neuroscience), Brain and Mind Centre, University of Sydney, NSW, Australia 2050.

luke.henderson@sydney.edu.au (email); +612 9351 7063 (Tel)

6. Number of Figures: 4

7. Number of Tables: 3

8. Number of Multimedia: 0

9. Number of words for Abstract: 230

10. Number of words for Significance Statement: 103

11. Number of words for Introduction: 499

12. Number of words for Discussion: 1700

13. Acknowledgements: The authors acknowledge the facilities and scientific and technical assistance of NeuRA Imaging, a node of the National Imaging Facility, a National Collaborative Research Infrastructure Strategy (NCRIS) capability.

14. Conflict of interest: The authors report no conflict of interest.

15. Funding: This research was supported by the National Health and Medical Research Council of Australia grant 1130280.

Abstract: Chronic neuropathic pain can result from nervous system injury and can persist in the absence of external stimuli. Although ongoing pain characterizes the disorder, in many individuals, the intensity of this ongoing pain fluctuates dramatically. Previously, it was identified that functional magnetic resonance imaging (fMRI) signal covariations between the midbrain periaqueductal gray matter (PAG), rostral ventromedial medulla (RVM), and spinal trigeminal nucleus (SpV) are associated with moment-to-moment fluctuations in pain intensity in individuals with painful trigeminal neuropathy (PTN). Since this brainstem circuit is modulated by higher brain input, we sought to determine which cortical sites might be influencing this brainstem network during spontaneous fluctuations in pain intensity. Over 12 minutes, we recorded ongoing pain intensity in 24 PTN participants, and classified them as fluctuating (n=13) or stable (n=11). Using a PAG seed, we identified connections between the PAG and emotional-affective sites such as the hippocampal and posterior cingulate cortices, the sensory-discriminative posterior insula, and cognitive-affective sites such as the dorsolateral prefrontal (dlPFC) and subgenual anterior cingulate cortices that were altered dependent on spontaneous high and low pain intensity. Additionally, sliding-window functional connectivity analysis revealed that the dlPFC-PAG connection anticorrelated with perceived pain intensity over the entire 12-minute period. These findings reveal cortical systems underlying moment-to-moment changes in perceived pain in PTN, which likely cause dysregulation in the brainstem circuits previously identified, and consequently alter the appraisal of pain across time.

Significance statement: Whilst the intensity of an individual's chronic pain is often measured at a specific point in time, it is known that in a large proportion of individuals, pain intensity fluctuates dramatically from moment-to-moment. In individuals with chronic neuropathic pain, we found that these spontaneous pain intensity fluctuations are associated with neural function fluctuations, specifically of function reflected as neural connectivity between brainstem pain modulatory circuits and cortical regions, including the dorsolateral prefrontal and cingulate cortices. These findings raise the possibility that modulating brain regions such as the dorsolateral prefrontal cortex in individuals with fluctuating chronic pain may provide an avenue for analgesic treatment.

Introduction:

The perception of pain is vital to survival, warning us of danger and motivating us to remove ourselves from it. As such, pain is adaptive. However, if pain persists beyond the period of normal healing and extends for longer than 3 months, it is considered chronic in nature, maladaptive, and pathological. Chronic pain that results from damage to the nervous system is called neuropathic pain and can occur in the absence of any external stimulus. In any one individual, the ongoing intensity of their pain can fluctuate between days, hours, or even from moment-to-moment. Whilst the precise biological mechanisms underlying perceived pain intensity fluctuations in chronic pain patients are unclear, it is well known that the brainstem contains multiple circuits capable of modulating the intensity of perceived pain by modulating noxious information at the level of the dorsal horn and spinal trigeminal nucleus (SpV) (Crawford et al, 2022b; Oliva et al, 2022; Youssef et al, 2016).

Previous work has demonstrated the role of these circuits in chronic neuropathic pain by establishing that signal coupling between the midbrain periaqueductal gray matter (PAG), rostral ventromedial medulla (RVM), and SpV was associated with perceived spontaneous pain intensity fluctuations in PTN patients (Mills et al, 2020). Whilst this circuit is a well-established brainstem pain modulatory network, whether changes in the connectivity of this circuit alone underpin spontaneous changes in pain intensity, or whether the connectivity of this brainstem circuit is also modulated by descending inputs from higher brain centres remains unknown. It is well-known that the PAG receives descending modulatory influences from higher brain centres including areas such as the dorsolateral prefrontal cortex (dlPFC), anterior cingulate cortex (ACC),

hippocampus, amygdala and the hypothalamus (Eippert et al, 2009; Ongür et al, 1998; Sevel et al, 2015).

These connections appear to drive changes in pain intensity in situations such as placebo and offset analgesia and nocebo hyperalgesia (Crawford et al, 2023a; Crawford et al, 2023b). Indeed, it was previously shown that signal covariation between these regions and the PAG are critical in determining whether an individual mounts a placebo analgesia (Crawford et al, 2023a). Further, it has also been shown that a connection between the dlPFC and PAG drives pain intensity variability during repeated noxious stimuli of the same intensity in healthy participants (Crawford et al, 2022a; Crawford et al, 2023b). Given these findings, it is likely that descending influences from higher brain regions also determine whether an individual displays spontaneous changes in ongoing pain intensity.

The aim of this investigation was to determine if signal covariations between higher cortical regions and the PAG are associated with moment-to-moment fluctuations in pain intensity in individuals with painful trigeminal neuropathy (PTN). We hypothesized that higher brain regions with well-established pain modulatory ability such as the dlPFC, ACC, hippocampus and amygdala would display increased functional connectivity with the PAG during periods of high relative to low pain intensity and would also display moment-to-moment functional connectivity changes with the PAG that were associated with fluctuations in perceived pain intensity.

Materials and methods:

Participants and Pain Measurements

Twenty-four PTN participants were recruited for the study (8 male; mean \pm SEM age=47.3 \pm 2.9 years; range=25-78 years). Diagnosis of PTN presence was made in accordance with the Liverpool criteria (Nurmikko & Eldridge, 2001) by a clinician in the research group (E.R.V). Inclusion criteria required a primary pain complaint relating to PTN and a pain intensity on the day of scanning >0. Exclusion criteria included typical MRI contraindications (current pregnancy, metallic implants), as well as the presence of any additional chronic pain condition in addition to PTN. Written informed consent was obtained from each participant for all procedures, approved by the [Author University] Human Research Ethics Committee and consistent with the declaration of Helsinki. Data from 24 PTN participants was used in previous studies (Mills et al, 2018; Mills et al, 2020).

For the seven days preceding the day of scanning, PTN participants recorded the intensity of their ongoing pain three times a day using a visual analogue scale (VAS) (0=no pain; 10=worst pain imaginable). The average of these pain ratings was taken as a measure of “diary pain intensity”. Participants also described their pain distribution by shading areas of a standard anatomical template of the face, and described the quality of their ongoing pain through completion of the McGill’s pain questionnaire (Melzack, 1975).

Scanning Procedures

Magnetic resonance imaging (MRI) was recorded during this study using a 3-Tesla MRI scanner (Achieva, Philips Medical Systems, The Netherlands) at [Author Scanning Location]. Participants lay supine on a scanner bed with their head positioned within a 32-channel transmit and receive head coil. Ear buds and over ear

headphones were provided to participants to reduce scanner noise, and sponges were placed between these headphones and the head coil to limit head motion throughout the course of functional scanning. A high-resolution T1-weighted anatomical image of the entire brain was collected (288 axial slices, repetition time=5600ms, raw voxel size=0.87x0.87x0.87mm thick, acquisition time=3min45sec). Following this, a series of 360 gradient-echo echo-planar brain volumes with blood oxygen level dependent (BOLD) contrast were recorded (37 axial slices, repetition time=2000ms, echo time=30ms, raw voxel size=3.0x3.0x4.0mm thick, acquisition time=12min). Throughout this 12-minute functional MRI (fMRI) scan, participants were instructed to rate their ongoing pain experience using a computerized visual analogue scale (CoVAS, Medoc, Israel). The CoVAS was connected to a digital screen visible to participants whilst inside the scanner, allowing them to report their ongoing pain intensity in real time (0=no pain; 10=worst pain imaginable). Pain intensity values were saved at the conclusion of each scanning session and subsequently averaged into 2-second rating periods overlying each functional volume for future analysis.

MRI and Statistical Analysis

Structural and functional image preprocessing was performed using Statistical Parametric Mapping version 12 (SPM12) within MATLAB version R2023a (Ashburner et al, 2014; Friston et al, 1994). fMRI images were first realigned and movement parameters inspected to ensure that no participant displayed excessive motion throughout the 12-minute scan (>1mm rotational or >0.5 degrees radian translational motion). These movement parameters were then modelled and removed from the fMRI signal through a Linear Modelling of Realignment Parameters (LMRP) procedure (Macey et al, 2004). Cardiac (frequency band of 60-120 beats per minute + 1

harmonic) and respiratory (frequency band of 8-25 breaths per minute + 1 harmonic) noise was modelled and removed using the Dynamic Retrospective Filtering (DRIFTER) toolbox (Särkkä et al, 2012). Each participant's fMRI image series was then linearly detrended to remove the effects of global signal intensity changes and co-registered to their corresponding T1-weighted anatomical image. The T1-weighted image was spatially normalized to the Montreal Neurological Institute (MNI) template using the Manual Computational Anatomy toolbox (CAT12) (Seiger et al, 2018) and these same normalization parameters applied to the fMRI image series. This process resulted in fMRI images being resliced into 2x2x2mm voxels. Finally, for the functional connectivity analyses, we applied a temporal high-pass filter with a cut-off frequency of 0.02Hz to the smoothed fMRI images to remove the low-frequency signals that cannot be resolved using 50-second windows (Leonardi & Van De Ville, 2015). Given that our elected seed resided within the midbrain PAG, a small structure with a discrete functional colocalization, all connectivity analyses were first performed using unsmoothed fMRI images to preserve the integrity of seed signal and seed-to-voxel relationships. Resulting connectivity contrast maps were then spatially smoothed using a 6mm Full Width at Half Maximum (FWHM) Gaussian filter.

Since equal number of PTN participants presented with unilateral left- and right-sided pain, functional images were flipped along the midline so that each participants resulting contrast images represented chronic pain in the left side of the face. Given that ascending pain pathways are largely represented contralaterally (at least in the case of the thalamus, the primary somatosensory and posterior insular cortices), this methodological consideration meant that we could produce a clear representation of

the cortical sites receiving input from, or driving the activity to the unilateral (right) PAG seed identified in previous investigations (Mills et al, 2020).

Pain Ratings

Inspection of pain intensity ratings across the entire 12-minutes of fMRI scanning revealed a stratification within PTN participants, such that in some individuals ongoing pain fluctuated greatly while in others pain remained relatively stable. In 11 participants, pain remained stable with these participants reporting an average difference between minimum and maximum pain of 0.2 out of 10 on the VAS (stable group). In the remaining 13 participants, ongoing pain intensity varied considerably, and in these participants, we observed an average difference in minimum to maximum pain of 3.4 out of 10 on the VAS (fluctuating group). For fMRI analyses, in order to match perceived pain intensity with changes in BOLD signal intensity and connectivity, ongoing pain intensity ratings for each participant were shifted forward two volumes (4 seconds). This shift accounted for the approximate hemodynamic delay since fMRI measures the oxygenation demands in discrete brain regions that follow 3-5 seconds after the neuronal response (West et al, 2019).

PAG Functional Connectivity:

Since a previous investigation (Mills et al, 2020) revealed the PAG, via relay with the RVM and SpV, to be an integral component of the descending brainstem circuitry responsible for spontaneous pain fluctuations in PTN (Mills et al, 2018) , we were interested in determining the cortical inputs which first make contact with this brainstem circuitry to drive these effects. As such, we performed functional connectivity using the PAG cluster previously identified in brainstem-isolated space as

a seed region for whole brain voxel-by-voxel functional connectivity analyses (Figure 1A). In the present study, the PAG seed comprised eight contiguous voxels, four voxels at two rostro caudal levels extending $z=-8$ to -10 in MNI space. The fMRI signal from each voxel within the seed was extracted and averaged to represent the mean PAG seed fMRI signal.

To study the functional cortical connections to the PAG, we performed separate functional connectivity analyses. We first calculated PAG connectivity within pre-defined time periods (blocks) corresponding to low and high pain intensities (Figure 1B). Secondly, we discerned additional temporal information about these connections by using a sliding-window dynamic functional connectivity analysis to assess the relationship between PAG connectivity and pain intensity throughout the 12-minute scan period (Figure 1C).

PAG High versus Low Pain Block Functional Connectivity

First, we aimed to determine how PAG connectivity with cortical sites was different in PTN participants during periods of spontaneously high relative to spontaneously low perceived pain. To achieve this, for each of the 13 fluctuating pain participants, we calculated PAG connectivity in two separated 50-second (25 volumes) time periods (blocks) – one during the lowest perceived pain period across the 12-minute fMRI scan and the other during the highest perceived pain (Figure 1B). Importantly, to minimize any motion-related artefact relating to moving the CoVAS slider, we isolated blocks where pain remained relatively stable for an entire 50-second period and avoided any periods when a large ($>1\text{cm}$) volume-to-volume movement along the CoVAS occurred. For each block, using each participant's fMRI images we conducted first-level

analyses with timeseries of PAG seed signal within the specified block attached as a condition, similar boxcar models traditionally used in task-related fMRI analyses. This first-level resulted in the generation of two contrast images for each participant, one where voxels of greater beta-value represented greater PAG connectivity during spontaneously low pain, and the other during spontaneously high pain. These images were spatially smoothed with a 6mm FWHM gaussian filter. These contrast images reflecting connectivity map effects were entered into a second level, paired, random-effects analysis of all 13 fluctuating pain participants. We set an initial threshold of $p < 0.001$ uncorrected for multiple comparisons, with a cluster defining threshold of 10 contiguous voxels, and each resulting cluster was subject to small volume correction (svc) for multiple comparisons to reduce the likelihood of type 1 errors.

Within each resulting cluster, we extracted PAG connectivity strength values for both lowest- and highest-pain blocks in each fluctuating pain participant and plotted mean (\pm SEM) functional connectivity strengths to provide a measure of connectivity direction (i.e. correlated or anti-correlated). These connectivity strengths were then entered into post-hoc paired t-tests to determine the direction and degree of significance in change of PAG connectivity between low- and high-pain blocks (Bonferroni corrected for multiple comparisons). In addition, because the majority of participants with fluctuating pain showed not one single VAS movement but several throughout the scan, which corresponded to pain intensity values which were between the lowest and highest blocks, we further assessed whether the sites we identified in the initial paired analysis also displayed a “dose response” with PAG connectivity and perceived pain. That is, additional blocks were included where possible where PAG signal was extracted, and contrast maps generated during 50-second periods of pain closer to the lowest

reported intensity (middle-low) or closer to the highest reported intensity (middle-high). This process resulted in a maximum of four PAG connectivity blocks for each cluster that represented fluctuating pain during lowest (block 1) to highest (block 4) pain in each participant.

To assess PAG connectivity over time in the 11 participants who displayed stable pain over the 12-minute scanning period, we calculated PAG connectivity in four equally spaced 50-second blocks (Figure 2A). For each significant cluster identified in our fluctuating pain participants low- versus high-pain block analysis, we extracted and plotted the mean PAG connectivity for all four blocks within the stable pain participants.

PAG Sliding-Window Functional Connectivity

Since it is likely that cortical structures engage pain modulatory circuits of the brainstem not only when pain is spontaneously high or low but also across a continuum, we determined if there was an ongoing relationship between fluctuating and spontaneous pain intensity and PAG connectivity strength. Eleven of the 13 fluctuating pain participants displayed multiple (>3) pain intensity changes throughout the 12-minute scanning period. Using the Dynamic BC toolbox (Liao et al, 2014), in each of these 11 participants, fMRI image sets were divided into 50-second (25 volume) windows with a repetition time of 2 seconds, resulting in 336 sliding windows (Figure 1C). For each of these windows, a Pearson's correlation coefficient was calculated between the PAG seed and each cortical voxel, resulting in a series of 336 dynamic PAG connectivity maps per participant. After undergoing spatial smoothing (6mm FWHM), these dynamic PAG connectivity maps were entered into a first-level linear regression analysis in which PAG dynamic functional connectivity was

compared with 336 corresponding sliding-window pain intensity ratings. This resulted in a single contrast image where more positive beta-values represented voxels where ongoing coupling with the PAG closely followed spontaneous pain fluctuations – that is, when voxel-PAG connectivity was higher, so too was perceived pain and vice versa for low pain.

These contrast images were then entered into a second-level one sample random effects analysis to determine cortical sites in which dynamic PAG functional connectivity significantly correlated with perceived pain in each of the 11 fluctuating pain participants. We set an initial threshold of $p < 0.001$ uncorrected for multiple comparisons, with a cluster defining threshold of 10 contiguous voxels and each resulting cluster was subject to small volume correction (svc) to reduce the likelihood of type 1 errors.

Results:

PTN Participant Characteristics

Individual PTN participant characteristics are displayed in Table 1. Mean (\pm SEM) pain in the 7-days preceding the MRI session (diary pain) was 3.8 ± 0.4 out of 10 with a mean PTN pain duration of 5.2 ± 1.3 years. All 24 PTN participants recorded current pain on the day of scanning, frequently selecting words to describe their pain such as throbbing (39%), sharp (26%), and exhausting (37%). Across the 12-minutes of scanning, 13 participants reported multiple fluctuations in pain intensity, with 11 participants reporting relatively stable pain intensity – altering VAS slider position a maximum of once within the entire 12-minute period (Figure 2A). Variable and consistent PTN participants did not differ significantly in terms of age ($mean \pm SEM$

years: Variable= 43.7 ± 4.0 ; Consistent= 51.5 ± 4.0 ; $p=0.20$), pain diary intensity ($mean \pm SEM$ pain: Variable = 4.3 ± 0.6 ; Consistent = 3.1 ± 0.4 ; $p=0.15$), or years lived with PTN pain ($mean \pm SEM$ years: Variable = 4.6 ± 1.8 ; Consistent = 5.9 ± 1.8 ; $p=0.66$) (two-sample t-test). Furthermore, the previous 7-days mean pain intensities were not predictive of pain during the scan which is revealed by the lack of correlation ($R = 0.22$) and no statistical significance ($p=0.33$) between these measures. Of the 13 variable PTN participants, 6 reported predominantly left-sided pain, and 7 predominantly right-sided pain. In the 11 consistent PTN participants, 3 reported predominantly right-sided pain, 5 left-sided pain, and 3 reported bilateral pain.

During the 12-minute fMRI scan, however, compared with PTN participants with consistent pain ratings those with variable ratings demonstrated significantly greater ongoing pain intensity ratings ($mean \pm SEM$ VAS pain: Variable = 3.9 ± 0.7 ; Consistent = 1.9 ± 0.6 ; $p<0.05$), as well as pain intensity variability ($mean \pm SEM$ VAS variability: Variable = 7.1 ± 1.3 ; Consistent = 0.4 ± 0.2 ; $p<0.05$) (two-sample t-tests) (Figure 2B). Despite this, after performing linear regression analyses in all 24 PTN participants, no significant interaction was observed between ongoing pain intensity and variability across the entire 12-minute scanning period ($R=0.27$, $p=0.20$) (Figure 2C). For the 13 fluctuating PTN participants, mean ($\pm SEM$) VAS pain intensity during the 4 pain blocks used for subsequent connectivity analyses were: Block 1 (lowest pain; $n=13$) 2.9 ± 0.7 , Block 2 ($n=11$) 3.7 ± 0.4 , Block 3 ($n=10$) 4.0 ± 0.8 , Block 4 (highest pain; $n=13$) 5.0 ± 0.7 .

PAG High versus Low Pain Block Functional Connectivity

Our voxel-by-voxel analysis comparing PAG functional connectivity during periods of spontaneously high versus low pain in our fluctuating PTN participants revealed

several cortical regions displaying pain intensity related changes. Significantly greater PAG connectivity strength during high relative to low pain was identified in the ipsilateral posterior insular cortex (mean±SEM PAG connectivity Beta-value lowest pain vs highest pain: 0.011 ± 0.009 vs 0.037 ± 0.013), ipsilateral hippocampus (-0.011 ± 0.005 vs 0.028 ± 0.010), ipsilateral dlPFC (-0.007 ± 0.004 vs 0.007 ± 0.003), and the posterior cingulate cortex (PCC) (-0.005 ± 0.004 vs 0.021 ± 0.006). The inverse contrast revealed a single cluster satisfying correction in the subgenual anterior cingulate cortex (sgACC) which displayed significantly greater PAG connectivity during periods of spontaneously low compared to high pain (0.023 ± 0.008 vs -0.001 ± 0.005) (Figure 3, Table 2) (paired t-tests, all $p<0.001$ uncorrected for multiple comparisons, small volume corrected). No significant linear relationship was observed between PAG connectivity within each of these clusters during low or high pain periods and either mean pain diary intensity or variability in the seven days preceding the scan (Table 1).

For each of these clusters, PAG connectivity strength was also extracted from additional blocks of increasing pain throughout the scan for a maximum of four graded pain blocks (block 2=moderate-low pain, block 3=moderate-high pain). Within the ipsilateral hippocampus (mean±SEM PAG connectivity Beta-value: block 2, 3: 0.007 ± 0.007 , 0.014 ± 0.009), dlPFC (0.005 ± 0.005 , 0.013 ± 0.008), and PCC (0.005 ± 0.005 , 0.013 ± 0.008) clusters, we observed a graded response of PAG connectivity change such that PAG connectivity closely followed spontaneous pain as it moved higher/lower throughout the scan. Conversely, clusters within the ipsilateral posterior insula and sgACC did not clearly show this graded response effect. Visual inspection of connectivity patterns within these clusters showed a strikingly similar

effect between each of our 13 fluctuating PTN participants, with 12 of the 13 participants showing increases between low and high pain periods in PAG-hippocampus, PAG-posterior insula, and PAG-dIPFC. Likewise, 12 of the 13 participants showed decreases between low and high pain periods in PAG-sgACC connectivity.

Within each of these five clusters, measures of PAG connectivity in stable pain PTN participants were extracted within four evenly distributed scan windows as described previously. In no single cluster did we identify changes in PAG connectivity throughout the scan duration. That is, within the 11 stable PTN participants, PAG connectivity with the ipsilateral posterior insula (mean \pm SEM PAG connectivity Beta-value Blocks 1, 2, 3, 4: 0.016 \pm 0.010, 0.006 \pm 0.012, 0.011 \pm 0.010, 0.016 \pm 0.010), the PCC (0.002 \pm 0.008, 0.004 \pm 0.008, 0.001 \pm 0.009, 0.005 \pm 0.008), the ipsilateral hippocampus (0.008 \pm 0.018, 0.009 \pm 0.014, 0.021 \pm 0.015, 0.013 \pm 0.013), the ipsilateral dIPFC (0.005 \pm 0.008, 0.010 \pm 0.006, 0.006 \pm 0.007, 0.010 \pm 0.004) nor the sgACC (-0.003 \pm 0.005, 0.005 \pm 0.007, 0.004 \pm 0.005, 0.006 \pm 0.006) were significantly altered between any of the four pain blocks (one-way ANOVA, $p>0.05$). Moreover, post-hoc two-factor repeated ANOVA analyses revealed a significant interaction between group assignment (i.e. “stable” or “fluctuating” PTN pain) and PAG connectivity strength with each of these five regions between each increasing intensity pain block (Figure 3-1).

PAG Sliding Window Connectivity

Dynamic functional connectivity analysis revealed that in our group of fluctuating PTN participants, PAG dynamic functional connectivity was inversely correlated with sliding-window pain intensity over the entire scan in a region of the contralateral (to

side of predominant pain) dlPFC (Figure 4, Table 3). We extracted and temporally smoothed the PAG dynamic connectivity values within the dlPFC cluster, identifying a significant inverse relationship between ongoing pain intensity ratings and PAG-dlPFC connectivity in 9 of the 11 participants ($p < 0.05$). That is, as pain was spontaneously high throughout the 12-minute scanning period, PAG-dlPFC connectivity was more anticorrelated, and vice versa. Two individual participant plots of pain versus PAG-dlPFC connectivity are shown in Figure 4. In no cortical region did PAG dynamic connectivity positively correlate with pain intensity.

Discussion:

The results of this study highlight that in individuals with chronic neuropathic pain, fluctuations in ongoing pain intensity are associated with altered communication between cortical areas with known roles in pain-processing and modulation, and pain-modulatory circuitry of the brainstem. That is, when pain was spontaneously high, we observed higher PAG coupling with the hippocampus, as well as the posterior insula, dlPFC, and PCC compared to periods where pain was spontaneously low. Conversely, when pain was spontaneously low, greater PAG coupling with the sgACC occurred compared to when pain was spontaneous high. These coupling changes were isolated to the fluctuating PTN participant group, with no overall changes in PAG coupling within these sites in PTN participants where pain remained stable. Additionally, our sliding-window analysis revealed that throughout the entire 12-minute scanning window, PAG-dlPFC coupling was negatively associated with ongoing pain intensity ratings. That is, the higher the pain the lower the PAG-dlPFC coupling and vice-versa. These data show that there are moment-to-moment changes in cortical signalling to, or reception of signals from, brainstem pain-modulatory circuits residing in the

midbrain PAG, and these likely contribute to fluctuations in spontaneous pain intensity in individuals suffering chronic neuropathic pain.

It is well-established that the PAG is capable of altering the perception of pain either by descending input to the spinal cord dorsal horn (for the body) or SpV (for the face) via relay with the RVM, or by ascending inputs to the cortical sites responsible for the pain percept (Fairhurst et al, 2007; Heinricher et al, 2009; Ong et al, 2019; Zambreanu et al, 2005). In pain-free individuals, the PAG-RVM descending pathway is believed finely balanced to manage nociceptive load, enabling analgesia when pain is unnecessary (such as when fighting or fleeing), or hyperalgesia to enable rest and healing behaviours. However, in individuals with chronic neuropathic pain, it has been suggested that this pathway is shifted, driven by cortical input, to promote an baseline pro-nociceptive state (Chen et al, 2018; Henderson & Keay, 2018; Ossipov et al, 2014). Indeed, it was previously established that this PAG-RVM circuitry was modulated in this group of PTN patients, with altered connectivity patterns between the PAG-RVM-SpV in those where pain fluctuated considerably over time. Here we establish the likely top-down cortical inputs which contact this network to drive pain intensity fluctuations.

It has been previously proposed that the dlPFC acts as a cortical interface between cognitive processing and pain modulation, performing this role by communicating with the PAG both directly and via the anterior cingulate cortex (Eippert et al, 2009; Sevel et al, 2015). Directed connectivity analyses previously identified decreases in dlPFC-PAG coupling as predictive of greater placebo analgesia in a healthy population (Sevel et al, 2015), and non-invasive stimulation of the dlPFC can both attenuate and amplify

endogenous pain modulatory responses (Krummenacher et al, 2010; Tu et al, 2021). Moreover, stimulation of the dlPFC in chronic pain patients can enhance pain tolerance and alleviate pain symptoms (Graff-Guerrero et al, 2005; Leung et al, 2018). Previously, the function of the dlPFC was extended to be involved in moment-to-moment pain intensity changes in healthy humans, with the dlPFC demonstrating reduced activation, alongside altered coupling patterns with pain processing areas such as the posterior insula and primary somatosensory cortex, in individuals who express greater variation in perceived pain intensity during a series of identical noxious stimuli (Crawford et al, 2022a). The results of the present investigation suggest that this dlPFC role is also operating in chronic pain patients - via descending input to the PAG - with both block and sliding window connectivity analyses showing dlPFC-PAG connectivity is associated with moment-to-moment fluctuations in pain intensity. Additionally, top-down connections from the dlPFC to PAG are known to not only encode pain modulation but also threat responses to an individual's immediate environment. Wang and colleagues demonstrated the functional connectivity patterns of distinct PAG columns during threat processing, showing that more negative PAG-dlPFC coupling during high threat relative to low threat states (Wang et al, 2022). Given that greater pain, especially when inside the confined environment of an MRI scanner, likely imposes a high threat to body homeostasis, ongoing and updating signals from the dlPFC to the PAG could encode such a threat.

Similar to the dlPFC, we identified that coupling between the PAG and posterior insula increased with perceived pain intensity. The posterior insular appears to encode sensory components of pain, and shows functional activation correlates with pain intensity in healthy humans (Wager et al, 2013). Uniquely, the posterior insular is one

of the only cortical sites where direct stimulation can trigger pain, and where cortical injury can cause the manifestation of neuropathic pain symptoms (Bergeron et al, 2021; Mazzola et al, 2012). Further, viral tract tracing in non-human primates demonstrates the posterior insula as the primary target for projection neurons arising from the spinothalamic tract (Dum et al, 2009), consistent with a role in detecting and encoding the magnitude of incoming pain signals. Since our fluctuating group displayed significantly higher variability of ongoing pain and greater overall pain intensity, the posterior insula may code both overall pain intensity and modulate the PAG to alter moment-to-moment pain intensity fluctuations.

It has been proposed that brain regions that code the emotional-affective aspects of pain, such as the hippocampus and posterior cingulate, play a more central role in chronic pain than they do during acute pain (Garcia-Larrea & Peyron, 2013). In humans, the connection between the hippocampus and PAG is believed to encode the memory and retrieval of pain and threat responses (Mokhtari et al, 2019; Wang et al, 2022), and in experimental animals, chemogenic modulation of hippocampal function can suppress neuropathic pain behaviours via direct modulation of sensory processing sites (Xuhong et al, 2018). Further, human chronic pain patients demonstrate increased ongoing activation in the PCC when compared with controls, and in fibromyalgia patients, this increased activation has shown to correlate with pain catastrophizing, or negative affect toward pain symptoms (Hsieh et al, 1995; Lee et al, 2018). Our data indicate that spontaneous changes in pain intensity in PTN patients are also encoded in these emotional-affective cortical regions, which may play a direct role in modulating the intensity of ongoing pain.

The sgACC was the only cortical region in which we found PAG connectivity to decrease as pain intensity spontaneously increased. The sgACC was originally proposed by Vogt and colleagues to rarely encode acute pain responses, and instead is integral for producing emotional responses to pain via direct autonomic projections to subcortical structures such as the central nucleus of the amygdala and the PAG (Vogt, 2005). These connections however have shown not only to encode emotional responses, but the degree of pain modulatory capacity in healthy humans. Specifically, in a large human cohort Cheng and colleagues showed that greater sgACC and brainstem coupling reduced the intensity of temporal summation (Cheng et al, 2015). Similarly, in a longitudinal study, Bingel and colleagues found that activation of the sgACC related to the magnitude of habituation over time, suggesting a core role of this site in suppressing nociceptive responses through recruitment of antinociceptive circuits (Bingel et al, 2008). Our data not only support the role of the sgACC in inhibiting pain responses over time, but further suggest that in PTN participants where pain fluctuates, the sgACC-PAG connection is inhibited to facilitate an increase in pain intensity.

Limitations: This study comes with several caveats. Firstly, we reflected some participants images so that the left side of the brain corresponded to the side of “most pain”. Whilst this approach was used to keep the pain side consistent, we appreciate that some higher brain regions display laterality and thus some information may have been diluted by this process. Secondly, since participants were required to operate the VAS manually by sliding their fingers along its extent, cortical activation of motor networks may have been induced that could alter our interpretation. To counter this, when conducting our pain block analysis, we isolated scan periods where pain

remained relatively stable within the 25-volume period of interest. Additionally, only the dIPFC survived correction in the dynamic functional connectivity analysis, which is not likely to be involved in directing hand movements and any motion effects would be averaged across the 50-second period comprising each of the 336-time windows. Thirdly, though our sample can detect a medium effect size with a power of 0.95, the relatively small sample size in both our fluctuating and stable pain groups could limit overall generalizability, although these sample sizes are similar to previous studies investigating ongoing pain fluctuations in chronic pain disorders (Baliki et al, 2011; Baliki et al, 2006; Foss et al, 2006; Geha et al, 2007). Finally, the choice to employ sliding-window connectivity over a “static connectivity” measure using each scan volume independently is worth noting. Whilst it is unclear whether averaging seed-to-voxel signal in short (50 second) periods measures similar processes observed across a total scan period, this technique has previously demonstrated utility in tying behavioural variables such as ongoing shifts in vigilance and respiration with physiological ones such as brain network dynamics (Chang et al, 2013; Thompson et al, 2013). Indeed, this investigation utilized a similar linear regression approach to Chang and colleagues, assessing whether PAG connectivity with the cortex tracks pain intensity over time.

Conclusions

Overall, our data reveal that cortical sites interact with the brainstem circuits responsible for ongoing pain intensity fluctuations in patients with chronic orofacial neuropathic pain. Whilst it is well known that areas like the dIPFC and ACC are involved in pain modulation in pain-free humans, we extend this idea and show that in individuals with ongoing pain, perceived intensity can fluctuate moment-to-moment

and we show here such fluctuation is associated with descending analgesic circuitry function. Given that areas such as the dIPFC are accessible to manipulation through techniques such as transcranial direct current stimulation (tDCS), transcranial alternating current stimulation (tACS), or repetitive transcranial magnetic stimulation (rTMS), these findings may provide the platform for testing the analgesic ability of the dIPFC in chronic pain, particularly in those individuals that display variations in ongoing pain intensity. Indeed, we have recently shown that tACS of the dIPFC can engage subcortical circuitry involved in the generation of sympathetic outflow to muscle and skin, likely via the PAG (Sesa-Ashton et al, 2022; Wong et al, 2023).

References:

- Ashburner, J., Barnes, G., Chen, C.-C., *et al* (2014) SPM12 manual. *Wellcome Trust Centre for Neuroimaging, London, UK*, 2464.
- Baliki, M. N., Baria, A. T. & Apkarian, A. V. (2011) The cortical rhythms of chronic back pain. *J Neurosci*, 31(39), 13981-90.
- Baliki, M. N., Chialvo, D. R., Geha, P. Y., *et al* (2006) Chronic pain and the emotional brain: specific brain activity associated with spontaneous fluctuations of intensity of chronic back pain. *J Neurosci*, 26(47), 12165-73.
- Bergeron, D., Obaid, S., Fournier-Gosselin, M. P., *et al* (2021) Deep Brain Stimulation of the Posterior Insula in Chronic Pain: A Theoretical Framework. *Brain Sci*, 11(5).
- Bingel, U., Herken, W., Teutsch, S., *et al* (2008) Habituation to painful stimulation involves the antinociceptive system - a 1-year follow-up of 10 participants. *PAIN*, 140(2), 393-394.
- Chang, C., Metzger, C. D., Glover, G. H., *et al* (2013) Association between heart rate variability and fluctuations in resting-state functional connectivity. *Neuroimage*, 68, 93-104.
- Chen, T., Taniguchi, W., Chen, Q.-Y., *et al* (2018) Top-down descending facilitation of spinal sensory excitatory transmission from the anterior cingulate cortex. *Nature Communications*, 9(1), 1886.
- Cheng, J. C., Erpelding, N., Kucyi, A., *et al* (2015) Individual Differences in Temporal Summation of Pain Reflect Pronociceptive and Antinociceptive Brain Structure and Function. *J Neurosci*, 35(26), 9689-700.
- Crawford, L., Mills, E., Meylakh, N., *et al* (2022a) Brain activity changes associated with pain perception variability. *Cerebral Cortex*.

571 Crawford, L. S., Boorman, D. C., Keay, K. A., *et al* (2022b) The pain conductor:
 572 brainstem modulation in acute and chronic pain. *Curr Opin Support Palliat Care*,
 573 16(2), 71-77.

574 Crawford, L. S., Meylakh, N., Macey, P. M., *et al* (2023a) Stimulus-independent and
 575 stimulus-dependent neural networks underpin placebo analgesia responsiveness in
 576 humans. *Communications Biology*, 6(1), 569.

577 Crawford, L. S., Mills, E. P., Peek, A., *et al* (2023b) Function and biochemistry of the
 578 dorsolateral prefrontal cortex during placebo analgesia: how the certainty of prior
 579 experiences shapes endogenous pain relief. *Cereb Cortex*, 33(17), 9822-9834.

580 Dum, R. P., Levinthal, D. J. & Strick, P. L. (2009) The spinothalamic system targets
 581 motor and sensory areas in the cerebral cortex of monkeys. *J Neurosci*, 29(45),
 582 14223-35.

583 Eippert, F., Bingel, U., Schoell, E. D., *et al* (2009) Activation of the Opioidergic
 584 Descending Pain Control System Underlies Placebo Analgesia. *Neuron*, 63(4), 533-
 585 543.

586 Fairhurst, M., Wiech, K., Dunckley, P., *et al* (2007) Anticipatory brainstem activity
 587 predicts neural processing of pain in humans. *Pain*, 128(12), 101-110.

588 Foss, J. M., Apkarian, A. V. & Chialvo, D. R. (2006) Dynamics of pain: fractal
 589 dimension of temporal variability of spontaneous pain differentiates between pain
 590 States. *J Neurophysiol*, 95(2), 730-6.

591 Friston, K. J., Holmes, A. P., Worsley, K. J., *et al* (1994) Statistical parametric maps
 592 in functional imaging: A general linear approach. *Human Brain Mapping*, 2(4), 189-
 593 210.

594 Garcia-Larrea, L. & Peyron, R. (2013) Pain matrices and neuropathic pain matrices:
 595 a review. *Pain*, 154 Suppl 1, S29-s43.

596 Geha, P. Y., Baliki, M. N., Chialvo, D. R., *et al* (2007) Brain activity for spontaneous
 597 pain of postherpetic neuralgia and its modulation by lidocaine patch therapy. *Pain*,
 598 128(1-2), 88-100.

599 Graff-Guerrero, A., González-Olvera, J., Fresán, A., *et al* (2005) Repetitive
 600 transcranial magnetic stimulation of dorsolateral prefrontal cortex increases
 601 tolerance to human experimental pain. *Brain Res Cogn Brain Res*, 25(1), 153-60.

602 Heinricher, M. M., Tavares, I., Leith, J. L., *et al* (2009) Descending control of
 603 nociception: Specificity, recruitment and plasticity. *Brain Res Rev*, 60(1), 214-25.

604 Henderson, L. A. & Keay, K. A. (2018) Imaging Acute and Chronic Pain in the
 605 Human Brainstem and Spinal Cord. *Neuroscientist*, 24(1), 84-96.

606 Hsieh, J.-C., Belfrage, M., Stone-Elander, S., *et al* (1995) Central representation of
 607 chronic ongoing neuropathic pain studied by positron emission tomography. *PAIN*,
 608 63(2), 225-236.

609 Krummenacher, P., Candia, V., Folkers, G., *et al* (2010) Prefrontal cortex modulates
 610 placebo analgesia. *PAIN*, 148(3), 368-374.

611 Lee, J., Protsenko, E., Lazaridiou, A., *et al* (2018) Posterior cingulate cortex is a key
 612 neural substrate for pain catastrophizing in fibromyalgia. *The Journal of Pain*, 19(3),
 613 S8.

614 Leonardi, N. & Van De Ville, D. (2015) On spurious and real fluctuations of dynamic
 615 functional connectivity during rest. *Neuroimage*, 104, 430-6.

616 Leung, A., Metzger-Smith, V., He, Y., *et al* (2018) Left Dorsolateral Prefrontal Cortex
 617 rTMS in Alleviating MTBI Related Headaches and Depressive Symptoms.
 618 *Neuromodulation*, 21(4), 390-401.

619 Liao, W., Wu, G.-R., Xu, Q., *et al* (2014) DynamicBC: a MATLAB toolbox for dynamic
 620 brain connectome analysis. *Brain connectivity*, 4(10), 780-790.

621 Macey, P. M., Macey, K. E., Kumar, R., *et al* (2004) A method for removal of global
622 effects from fMRI time series. *Neuroimage*, 22(1), 360-6.

623 Mazzola, L., Isnard, J., Peyron, R., *et al* (2012) Stimulation of the human cortex and
624 the experience of pain: Wilder Penfield's observations revisited. *Brain*, 135(Pt 2),
625 631-40.

626 Melzack, R. (1975) The McGill Pain Questionnaire: Major properties and scoring
627 methods. *Pain*, 1(3), 277-299.

628 Mills, E. P., Alshelh, Z., Kosanovic, D., *et al* (2020) Altered Brainstem Pain-
629 Modulation Circuitry Connectivity During Spontaneous Pain Intensity Fluctuations. *J*
630 *Pain Res*, 13, 2223-2235.

631 Mills, E. P., Pietro, F. D., Alshelh, Z., *et al* (2018) Brainstem Pain-Control Circuitry
632 Connectivity in Chronic Neuropathic Pain. *The Journal of Neuroscience*, 38(2), 465-
633 473.

634 Mokhtari, T., Tu, Y. & Hu, L. (2019) Involvement of the hippocampus in chronic pain
635 and depression. *Brain Science Advances*, 5(4), 288-298.

636 Nurmikko, T. J. & Eldridge, P. R. (2001) Trigeminal neuralgia--pathophysiology,
637 diagnosis and current treatment. *Br J Anaesth*, 87(1), 117-32.

638 Oliva, V., Hartley-Davies, R., Moran, R., *et al* (2022) Simultaneous brain, brainstem,
639 and spinal cord pharmacological-fMRI reveals involvement of an endogenous opioid
640 network in attentional analgesia. *Elife*, 11.

641 Ong, W.-Y., Stohler, C. S. & Herr, D. R. (2019) Role of the Prefrontal Cortex in Pain
642 Processing. *Molecular neurobiology*, 56(2), 1137-1166.

643 Ongür, D., An, X. & Price, J. L. (1998) Prefrontal cortical projections to the
644 hypothalamus in macaque monkeys. *J Comp Neurol*, 401(4), 480-505.

645 Ossipov, M. H., Morimura, K. & Porreca, F. (2014) Descending pain modulation and
646 chronification of pain. *Current opinion in supportive and palliative care*, 8(2), 143-
647 151.

648 Särkkä, S., Solin, A., Nummenmaa, A., *et al* (2012) Dynamic retrospective filtering of
649 physiological noise in BOLD fMRI: DRIFTER. *NeuroImage*, 60, 1517-27.

650 Seiger, R., Ganger, S., Kranz, G. S., *et al* (2018) Cortical thickness estimations of
651 FreeSurfer and the CAT12 toolbox in patients with Alzheimer's disease and healthy
652 controls. *Journal of Neuroimaging*, 28(5), 515-523.

653 Sesa-Ashton, G., Wong, R., McCarthy, B., *et al* (2022) Stimulation of the dorsolateral
654 prefrontal cortex modulates muscle sympathetic nerve activity and blood pressure in
655 humans. *Cereb Cortex Commun*, 3(2), tgac017.

656 Sevel, L. S., Craggs, J. G., Price, D. D., *et al* (2015) Placebo Analgesia Enhances
657 Descending Pain-Related Effective Connectivity: A Dynamic Causal Modeling Study
658 of Endogenous Pain Modulation. *The Journal of Pain*, 16(8), 760-768.

659 Thompson, G. J., Magnuson, M. E., Merritt, M. D., *et al* (2013) Short-time windows of
660 correlation between large-scale functional brain networks predict vigilance
661 intraindividually and interindividually. *Hum Brain Mapp*, 34(12), 3280-98.

662 Tu, Y., Wilson, G., Camprodon, J., *et al* (2021) Manipulating placebo analgesia and
663 nocebo hyperalgesia by changing brain excitability. *Proc Natl Acad Sci U S A*,
664 118(19).

665 Vogt, B. A. (2005) Pain and emotion interactions in subregions of the cingulate
666 gyrus. *Nature reviews. Neuroscience*, 6(7), 533-544.

667 Wager, T. D., Atlas, L. Y., Lindquist, M. A., *et al* (2013) An fMRI-based neurologic
668 signature of physical pain. *N Engl J Med*, 368(15), 1388-97.

Wang, S., Veinot, J., Goyal, A., *et al* (2022) Distinct networks of periaqueductal gray columns in pain and threat processing. *Neuroimage*, 250, 118936.

West, K. L., Zuppichini, M. D., Turner, M. P., *et al* (2019) BOLD hemodynamic response function changes significantly with healthy aging. *Neuroimage*, 188, 198-207.

Wong, R., Sesa-Ashton, G., Datta, S., *et al* (2023) The role of the dorsolateral prefrontal cortex in control of skin sympathetic nerve activity in humans. *Cereb Cortex*, 33(13), 8265-8272.

Xuhong, W., Ren, W., Centeno, M. V., *et al* (2018) Dorsal Hippocampal Activation Suppresses Neuropathic Pain Behaviors: Chronic pain as extinction-resistant pain-related memory traces. *bioRxiv*, 292094.

Youssef, A. M., Macefield, V. G. & Henderson, L. A. (2016) Pain inhibits pain; human brainstem mechanisms. *Neuroimage*, 124(Pt A), 54-62.

Zambreanu, L., Wise, R. G., Brooks, J. C. W., *et al* (2005) A role for the brainstem in central sensitisation in humans. Evidence from functional magnetic resonance imaging. *PAIN*, 114(3).

Figure Legends:

Figure 1. Midbrain periaqueductal gray (PAG) seed region, pain block, and sliding window connectivity design. **A.** With coordinates derived from a previous investigation utilizing this same data set, a 1mm sphere surrounding the PAG was generated and used as a seed region for subsequent connectivity analyses. **B.** Visual representation of the block analysis used to assess cortico-brainstem functional connectivity during periods of spontaneously high versus spontaneously low pain. In this analysis, connectivity between the PAG seed and each cortical voxel was assessed within a 50-second period overlying participants lowest reported pain (blue shaded bar) against a 50-second period overlying participants' highest reported pain (orange shaded bar). Pain intensity was assessed using a visual analogue scale (VAS) from 0=no pain to 10=worst pain imaginable. **C.** In addition to the block design, a sliding window analysis was performed where PAG connectivity was assessed during a 50-second period (window 1 light grey box) which was then shifted forward by one volume (window 2 dark grey box). This resulted in 336 contrast images representing 50-second mean PAG connectivity values across the entire 12-minute scan.

Figure 2. Ongoing pain intensity ratings and pain intensity variability in fluctuating (n=13) and stable (n=11) pain groups. **A.** Trace of VAS responses recorded throughout the entire 12-minute fMRI scan series. All participants fit the diagnostic criteria for trigeminal neuropathic pain, however, note that in only our fluctuating pain group (left plot) are multiple spontaneous shifts in perceived pain observed. **B.** Mean pain intensity and pain rating variability in fluctuating (orange bars) and stable (blue bars) pain groups across the 12-minute scan. Fluctuating pain participants exhibited significantly higher mean intensity and variability of pain than

stable participants. **C.** Linear regression analysis comparing participant mean pain intensity and variability throughout the 12-minute scan period. No significant interaction between these variables was observed despite variable participants demonstrating both greater intensity and variability than stable pain participants.

Figure 3. Significant differences in PAG connectivity strength during periods of high- versus low-pain in fluctuating pain participants (n=13). Across the 12-minute scanning window, the 25 volumes where participants rated their pain as most intense were entered into a second-level, paired design with the 25 volumes where participants rated their pain the least intense. The upper panel shows significant cortical clusters with greater midbrain periaqueductal gray matter (PAG) connectivity during high versus low pain (red colour scale) and vice versa (blue colour scale). The slice locations in Montreal Neurological Institute space are shown at the top right of each slice. The middle panel shows plots of mean (\pm SEM) PAG connectivity extracted from the above clusters and compared between fluctuating (high- and low-blocks) and stable (evenly dispersed blocks) participants. Note that the magnitude of PAG connectivity change in these sites only differs in those participants where pain fluctuates. Within the bottom panel is a post-hoc analysis of the dose-dependent response within these clusters, as across the scan course our fluctuating pain group exhibited periods of perceived pain between the lowest and highest reported blocks. Note that in the hippocampus, dorsolateral prefrontal, and posterior cingulate cortex PAG connectivity appears to particularly track with the magnitude of pain only in those participants displaying fluctuating pain. Please see Figure 3-1.

Figure 4. Cortical region demonstrating ongoing PAG connectivity change relating to fluctuations in pain intensity perceived across the 12-minute scan period. The left overlay shows firstly the midbrain periaqueductal gray matter (PAG) PAG seed (upper box) sectioned within a render also showing a significant cluster in the contralateral (to side of pain) dorsolateral prefrontal cortex where PAG-connectivity was anticorrelated with perceived pain throughout the entire scanning period in fluctuating pain patients. The right panel provides visual examples of two fluctuating pain PTN participants where PAG-dorsolateral prefrontal cortex connectivity (orange line) was continuously anticorrelated with perceived pain (blue line) over the 12-minute scanning window. FC = functional connectivity; VAS = visual analogue scale; dlPFC = dorsolateral prefrontal cortex.

Table Legends:

Table 1. Painful Trigeminal Neuropathy Participant Classification. Participants 1-13 were placed in the fluctuating pain group, and participants 14-23 in the stable pain group. *PEA = Palmitoylethanolamide; VAS = visual analogue scale; SD = standard deviation.*

Table 2. Montreal Neurological Institute co-ordinates, cluster sizes, and significance values for clusters showing midbrain periaqueductal gray matter connectivity differences during spontaneous low- vs high-pain intensity blocks in fluctuating pain participants (n=13).

Table 3. Montreal Neurological Institute co-ordinates, cluster sizes, and significance values for clusters showing correlations pain intensity and PAG connectivity across the entire 12-minute scan length in fluctuating pain participants (n=13).

eNeuro Accepted Manuscript

Table 1. Painful Trigeminal Neuropathy Participant Classification. Participants 1-13 were placed in the fluctuating pain group, and participants 14-23 in the stable pain group. *PEA = Palmitoylethanolamide; VAS = visual analogue scale; SD = standard deviation.*

	Age (Years)	Sex	Side of Pain	Diary Pain Intensity (VAS)	Diary Pain Variability (SD)	Pain Duration (Months)	Current medication Use
1	29	F	Left	5.3	1.5	49	-
2	45	M	Left	3.1	0.4	21	-
3	35	M	Right	1.5	0.7	96	PEA
4	78	F	Left	1.9	0.7	314	Paracetamol, Ostelin, Pritiq, Somac
5	47	F	Right	2.1	0.4	13	PEA
6	34	M	Right	3.7	1.7	10	PEA, Mersyndol
7	66	M	Left	3.9	0.6	48	-
8	44	F	Right	-	-	5	Valium, Panadine forte
9	47	F	Right	4.1	1.6	7	-
10	44	F	Left	8.1	0.4	75	PEA
11	46	F	Left	7.5	2.5	27	Endep
12	28	F	Right	5.8	0.3	52	-
13	25	M	Right	4.5	1.7	8	Panadine forte

14	69	F	Right	3.9	0.4	259	Endep, Lyrica, Oroxine, Crestor, Effexor, Folic acid, Prednisone
15	45	F	Bilateral	2.8	2.8	24	Duloxetine
16	66	M	Right	1.8	0.7	81	Endep
17	36	F	Left	1.9	0.6	37	PEA
18	40	F	Left	5.8	0.3	33	-
19	58	F	Left	4.0	0.4	11	Namipril, Zoloft, Panadol, PEA
20	35	M	Left	4.2	0.7	60	-
21	67	M	Bilateral	2.0	0.9	96	Endep
22	51	F	Left	1.5	0.3	13	Mobic, PEA
23	35	F	Right	3.4	1.0	15	Amitriptyline
24	64	F	Bilateral	-	-	144	-

767

768

769

Table 2. Montreal Neurological Institute co-ordinates, cluster sizes, and significance values for clusters showing midbrain periaqueductal gray matter connectivity differences during spontaneous low- vs high-pain intensity blocks in fluctuating pain participants (n=13).

	MNI Coordinate					
Region	X	Y	Z	Cluster size (kE)	t-value	z-value
<i>highest > lowest Pain</i>						
ipsilateral posterior insula	-36	-32	8	21	6.08	4.03
posterior cingulate cortex	-6	-36	30	18	5.16	3.68
ipsilateral hippocampus	-52	0	2	11	3.86	3.05
ipsilateral dorsolateral prefrontal cortex	-20	34	28	11	3.77	3.00
<i>Highest < Lowest Pain</i>						
subgenual anterior cingulate cortex	6	-28	-6	12	3.86	3.05

Table 3. Montreal Neurological Institute co-ordinates, cluster sizes, and significance values for clusters showing correlations pain intensity and PAG connectivity across the entire 12-minute scan length in fluctuating pain participants (n=13).

	MNI Coordinate					
Region	X	Y	Z	Cluster size (kE)	t-value	z-value
Negative Correlation						
contralateral dorsolateral prefrontal cortex	32	40	16	21	4.55	3.27







